

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
31 May 2001 (31.05.2001)

PCT

(10) International Publication Number  
**WO 01/37869 A1**

(51) International Patent Classification<sup>7</sup>: **A61K 39/385,**  
39/29, A61P 31/12

Corporation, 4560 Horton Street - R440, Emeryville, CA  
94608 (US).

(21) International Application Number: PCT/AU00/01410

(74) Agents: **SLATTERY, John, M. et al.;** Davies Collison  
Cave, 1 Little Collins Street, Melbourne, VIC 3000 (AU).

(22) International Filing Date:

17 November 2000 (17.11.2000)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

60/166,652 19 November 1999 (19.11.1999) US  
60/224,362 11 August 2000 (11.08.2000) US

(71) Applicants: **CSL LIMITED** [AU/AU]; 45 Poplar Avenue,  
Parkville, VIC 3052 (AU). **CHIRON CORPORATION**  
[US/US]; 4560 Horton Street, Emeryville, CA 94608-2916  
(US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,  
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,  
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,  
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,  
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,  
TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian  
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European  
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,  
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,  
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— With international search report.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(72) Inventors: **DRANE, Debbie;** 72 Mulcahy Road, Bullengarook, VIC 3437 (AU). **COX, John;** 140 Dunns Road, Bullengarook, VIC 3437 (AU). **HOUGHTON, Michael;** Chiron Corporation, 4560 Horton Street - R440, Emeryville, CA 94608 (US). **PALIARD, Xavier;** Chiron

(54) Title: VACCINE COMPOSITIONS

(57) Abstract: The present invention relates generally to an immunogenic complex comprising a charged organic carrier and a charged antigen and, more particularly, a negatively charged organic carrier and a positively charged antigen, wherein the charged antigen is a polypeptide of Hepatitis C Virus (HCV), particularly the core protein of HCV, or a fragment thereof, or a fusion protein comprising the polypeptide or a fragment thereof. The complexes of the present invention are useful, *inter alia*, in vaccine compositions as therapeutic and/or prophylactic agents for facilitating the induction of immune responses, and in particular a cytotoxic T-lymphocyte response, in the treatment of a disease condition which results from an HCV infection.

WO 01/37869 A1

B1

## VACCINE COMPOSITIONS

### 5 FIELD OF THE INVENTION

The present invention relates generally to a vaccine composition and to an immunogenic complex for use in the vaccine composition. In particular, the invention relates to an immunogenic complex comprising a charged organic carrier, more particularly a negatively charged organic carrier, and a charged antigen, more particularly a positively charged antigen, wherein the charged antigen is a polypeptide, preferably the core protein, of Hepatitis C Virus (HCV) or a fragment thereof, or a fusion protein comprising said polypeptide or a fragment thereof. The vaccine compositions and immunogenic complexes of the present invention are useful, *inter alia*, as therapeutic and/or prophylactic agents for facilitating the induction of immune responses, and in particular a cytotoxic T-lymphocyte response, in the treatment of a disease condition which results from an HCV infection.

### BACKGROUND OF THE INVENTION

Hepatitis C virus (HCV) is now recognised as the leading cause of chronic liver disease and an estimated 170 million people are currently infected with the virus (Armstrong, G. L., M. J. Alter, G. M. McQuillan, and H. S. Margolis. 2000. *Hepatology* 31:777, Cohen, J. 1999. *Science* 285:26). Despite these alarming numbers, very few therapies are available for treatment and those available are of low efficacy. Improved therapies are desperately needed but their development has been hampered by the lack of a small animal model for infection and disease, and inefficient systems for cultivating the virus. Several studies have suggested that T cell-mediated immune responses to HCV infection can affect the outcome of HCV infection and disease (Missale, G., R. Bertoni, V. Lamonaca, A. Valli, M. Massari, C. Mori, M. G. Rumi, M. Houghton, F. Fiaccadori, and C. Ferrari. 1996. *J. Clin. Invest.* 98:706., Cooper, S. L., A. L. Erickson, E. J. Adams, J. Kansopon, A. J. Weiner, D.

- 2 -

Y. Chien, M. Houghton, P. Parham, and C. M. Walker. 1999. *Immunity* 10:439, Lechner, F., D. K. Wong, P. R. Dunbar, R. Chapman, R. T. Chung, P. Dohrenwend, G. Robbins, R. Phillips, P. Klenerman, and B. D. Walker. 2000. *J. Exp. Med.* 191:1499). Furthermore there is an inverse correlation between the frequency of  
5 HCV specific cytotoxic T cells (CTLs) and viral load and the presence of HCV Core specific CTLs prior to interferon treatment has been associated with subsequent response of patients to this therapy (Nelson, D., C. Marousis, G. Davis, C. Rice, J. Wong, M. Houghton, and J. Lau. 1997. *J. Immunol.* 158:1473. Nelson, D. R., C. G. Marousis, T. Ohno, G. L. Davis, and J. Y. Lau. 1998. *Hepatology* 28:225 ). Thus, a  
10 vaccine capable of eliciting CTLs may be useful in control of HCV particularly as an adjunct to current therapies. The use of HCV proteins for vaccine development has been previously described (Houghton, EP Patent No. 0318216).

The adjuvant properties of saponin have been long known, as has its ability to  
15 increase antibody titres to immunogens. As used herein, the term "saponin" refers to a group of surface-active glycosides of plant origin composed of a hydrophilic region (usually several sugar chains) in association with a hydrophobic region of either steroid or triterpenoid structure. Although saponin is available from a number of diverse sources, saponins with useful adjuvant activity have been derived from the  
20 South American tree *Quillaja saponaria* (Molina). Saponin from this source was used to isolate a "homogeneous" fraction denoted "Quil A" (Dalsgaard, K., (1974), *Arch. Gesamte Virusforsch.* 44:243).

Dose-site reactivity is a major concern for both the veterinary and human use of Quil  
25 A in vaccine preparations. One way to avoid this toxicity of Quil A is the use of an immunostimulating complex (known as an ISCOM™, an abbreviation for Immuno Stimulating COMplex). This is primarily because Quil A is less reactive when incorporated into immunostimulating complexes, because its association with cholesterol in the complex reduces its ability to bind to cholesterol in cell membranes  
30 and hence its cell lytic effects. In addition, a lesser amount of Quil A is required to generate a similar level of adjuvant effect.

- 3 -

The immunomodulatory properties of the Quil A saponins and the additional benefits to be derived from these saponins when they are incorporated into an immunostimulating complex have been described in various publications, e.g. Cox and Cox, J.C. and Coulter, A.R. *Advances in Adjuvant Technology and Application* 5 *in Animal Parasite Control Utilising Biotechnology*, Chapter 4, Editor Yong, W.K. CRC Press (1992); Cox, J.C. and Coulter, A.R. (1997) *Vaccine*, 15(3):248-256; Cox, J.C. and Coulter, A.R. (1999) *BioDrugs* 12(6):439-453; Dalsgaard, (1974) (supra); Morein *et al.*, (1989) "Immunostimulating complex (ISCOM)", *In "Vaccines: Recent Trends and Progress"*. G. Gregoriadis, A.C. Allison and G. Poster (Eds). Plenum 10 Press, New York, p.153; Australian Patent Specifications Nos. 558258, 589915, 590904 and 632067.

Classic ISCOMs are formed by combination of cholesterol, saponin, phospholipid, and immunogens, such as viral envelope proteins. ISCOM matrix compositions 15 (known as ISCOMATRIX™) are formed identically, but without viral proteins. ISCOMs appear to stimulate both humoral and cellular immune responses. ISCOMs have been made with proteins from various viruses, including HSV-1, CMV, EBV, hepatitis B virus (HBV), rabies virus, and influenza virus, see for example, I.G. Barr *et al.*, *Adv. Drug Delivery Reviews*, 32:247-271 (1998). It has been observed that 20 where naked DNA or polypeptides from infectious agents are poorly immunogenic when given by themselves, inclusion within ISCOMs has increased their immunogenicity. Various proteins formulated with ISCOMs have been shown to induce CTL, mainly in rodent models. Berzofsky, (1991), *Biotechnol. Ther.* 2:123-135; Hsu *et al.*, (1996), *Vaccine* 14:1159-1166; Lipford *et al.*, (1994), *Vaccine* 12:73-80; 25 Mowat *et al.*, (1991), *Immunology* 72:317-322; Osterhaus *et al.*, (1998), *Dev. Biol. Stand.* 92:49-58; Rimmelzwaan *et al.*, (1997), *J. Gen. Virol.* 78 (pt.4):757-765; Sambhara *et al.*, (1998), *J. Infect. Dis.* 177:1266-1274; Sambhara *et al.*, (1997), *Mech. Aging Dev.* 96:157-169; Sjolander *et al.*, (1997), *Vaccine* 15:1030-1038; Sjolander *et al.*, (1998), *J. Leukoc. Biol.* 64:713-723; Takahashi *et al.*, (1990), 30 *Nature* 344:873-875; Tarpey *et al.*, (1996), *Vaccine* 14:230-236; Trudel *et al.*, (1987), *Vaccine* 10:107-112; Verschoor *et al.*, (1999), *J. Virol.* 73:3292-3300;

Villacres-Eriksson, (1995), *Clin. Exp. Immunol.* **102**:46-52; Zugel et al., (1995), *Eur. J. Immunol.* **25**:451-458.

Association between antigen and adjuvant is thought to be important for optimal  
5 induction of immune responses in particular CTL (Cox, J.C. and Coulter, A.R. (1999)  
*BioDrugs*, **12**(6):439-445). A number of studies have been done which confirm this  
hypothesis including work with virosomes and ISCOMs™ (Ennis, F. A., Crux, J.,  
Jameson, J., Klein, M., Burt, D. and Thipphawong, J. 1999. *Virology* **259**: 256-261.,  
Zurbriggen, R., Novak-Hofer, I., Seelig, A. and Gluck, R. (2000), *Progress in lipid*  
10 *Research* **39**: 3-18., Voeten, J. T. M., Nieuwkoop, N. J., Lövgren-Bengtsson, K.,  
Osterhaus, D. M. E. and Rimmelzwaan, G. F. 2000. *Euro J Imm (Submitted)*).  
Typically association between ISCOM™ and antigen has been achieved by  
incorporation of amphipathic antigens into the ISCOM™ structure during formation  
(Morein, B., B. Sundquist, S. Hoglund, K. Dalsgaard, and A. Osterhaus. 1984.  
15 *Nature* **308**:457). Incorporation was by hydrophobic interactions. However many  
antigens, such as the HCV core protein, were unable to be incorporated into  
ISCOMs™ by the classical method. More recently methods to associate antigens with  
a preformed protein-free immunostimulating complex (ISCOMATRIX™) utilising  
chelating and electrostatic interactions have been developed (International Patent  
20 Applications Nos. PCT/AU98/00080 – WO 98/36772, and PCT/AU00/00110).

The Core protein of HCV, as well as the E1 and E2 envelope proteins, have been  
shown to be useful in immunizing against HCV (see, e.g., copending U.S. Patent  
Application Serial No. 08/823,980). The sequences for the envelope proteins also  
25 contain certain conserved regions, even in the hypervariable domains thereof, which  
provides increased utility for immunization against the various escape mutants  
responsible for chronic infections. There remains a need, however, for methods and  
compositions which increase the ability of these proteins and polypeptides to be  
immunogenic and to stimulate the development of broad and durable CTL active  
30 against HCV infection.

- 5 -

In work leading up to the present invention, the inventors have developed an immunogenic complex based on the electrostatic association of an antigen of HCV and an organic carrier, such as an adjuvant. This electrostatic association permits co-delivery of the antigen and the organic carrier to the immune system, for the purpose of inducing an immune response, particularly a cytotoxic T-lymphocyte response, to the antigen.

---

#### SUMMARY OF THE INVENTION

Throughout this specification and the claims which follow, unless the context requires otherwise, the word "comprise", and variations such as "comprises" and "comprising", will be understood to imply the inclusion of a stated integer or step or group of integers or steps but not the exclusion of any other integer or step or group of integers or steps.

15

One aspect of the present invention relates to an immunogenic complex comprising a charged organic carrier and a charged antigen, which organic carrier and antigen are electrostatically associated, and wherein the charged antigen is a polyprotein of Hepatitis C Virus (HCV) or a fragment thereof, or a fusion protein comprising said polyprotein or a fragment thereof.

20

Preferably, the polyprotein is the core protein of HCV.

Another aspect of the present invention more particularly provides an immunogenic complex as described above, wherein the charged organic carrier is a negatively charged organic carrier.

25

Yet another aspect of the present invention provides an immunogenic complex as described above, wherein the charged organic carrier is a negatively charged adjuvant.

30

- 6 -

Yet still another aspect of the present invention provides an immunogenic complex as described above, wherein said negatively charged adjuvant is a naturally negatively charged adjuvant which has been modified to increase the degree of its negative charge.

5

A further aspect of the present invention relates to a vaccine composition comprising as the active component an immunogenic complex comprising a charged organic carrier and a charged antigen, which organic carrier and antigen are electrostatically associated, and wherein the charged antigen is a polyprotein of Hepatitis C Virus (HCV) or a fragment thereof, or a fusion protein comprising said polyprotein or a  
10 fragment thereof, together with one or more pharmaceutically acceptable carriers and/or diluents.

Preferably, the polyprotein is the core protein of HCV.

15

Another further aspect of the present invention relates to a method of eliciting, inducing or otherwise facilitating, in a mammal, an immune response to an antigen, said method comprising administering to said mammal an effective amount of an immunogenic complex or a vaccine composition as hereinbefore described.

20

Yet another further aspect of the present invention relates to a method of treating a disease condition in a mammal, said method comprising administering to said mammal an effective amount of an immunogenic complex or a vaccine composition as hereinbefore described, wherein administering said composition elicits, induces  
25 or otherwise facilitates an immune response which inhibits, halts, delays or prevents the onset or progression of the disease condition.

Still another further aspect the present invention relates to the use an immunogenic complex or vaccine composition as hereinbefore described in the manufacture of a  
30 medicament for inhibiting, halting, delaying or preventing the onset or progression of a disease condition.

- 7 -

Still yet another further aspect of the present invention relates to an agent for use in inhibiting, halting, delaying or preventing the onset or progression of a disease condition, said agent comprising an immunogenic complex or vaccine composition as hereinbefore described.

5

As described further below, the immunogenic complexes of the present invention may include, as the charged antigen associated with the charged organic carrier, an HCV protein such as an HCV Core nucleocapsid protein, a nonstructural protein, the E1 envelope protein, the E2 envelope protein, immunogenic fragments of any of such  
10 proteins, or combinations of such proteins. Such fragments generally include polypeptides comprising epitopes recognizable by T cells. Preferred fragments comprise those fragments which are immunogenic when provided by themselves, or when included in an immunogenic complex of the present invention. As used throughout this specification, the term "polyprotein of HCV" or "HCV protein" is  
15 intended to include the full length protein as well as polypeptide fragments. The HCV protein may also be present in the immunogenic complexes of the present invention as fusion proteins, depending on which method of expression of the HCV protein is chosen. The sequences for these polypeptides and proteins are known (see, e.g., U.S. Patent No. 5,350,671). The invention also provides polypeptides that are  
20 homologous (i.e., have sequence identity) to these fragments. Depending on the particular fragment, the degree of sequence identity is preferably greater than 50% (e.g., 60%, 70%, 80%, 90%, 95%, 99% or more). These homologous polypeptides include mutants and allelic variants of the fragments.

25

## BRIEF DESCRIPTION OF THE DRAWINGS

**Figure 1** is a graphical representation of the sucrose gradient analysis of Core-ISCOTM formulations for core protein (Figure 1A), ISCOMATRIXTM (Figure 1B), and  
30 Core-ISCOTM (Figure 1C).

**Figure 2:** B-LCLs from the two non-responders, HCV Core-ISCOTM-immunized



- 8 -

animals cannot present Core-derived peptides 121-135 and 86-100. Figure 2A: The 121-135-specific CTL line, established from animal DV037 was tested in a standard  $^{51}\text{Cr}$  release assay for its ability to lyse DV037 B-LCL target cells sensitized with peptide 121-135 (solid squares) or an irrelevant peptide (open squares), AY921 B-LCL target cells sensitized with peptide 121-135 (solid squares) or an irrelevant peptide (open circles), and BB231 B-LCL target cells sensitized with peptide 121-135 (solid triangles) or an irrelevant peptide (open triangles). Figure 2B: The 86-100-specific CTL line was tested in a standard  $^{51}\text{Cr}$  release assay for its ability to lyse DV036 B-LCL target cells sensitized with peptide 86-100 (solid squares) or an irrelevant peptide (open squares), AY921 B-LCL target cells sensitized with peptide 86-100 (solid circles) or an irrelevant peptide (open circles), and BB231 B-LCL target cells sensitized with peptide 86-100 (solid triangles) or an irrelevant peptide (open triangles).

Figure 3 shows the longevity of the CTL responses primed by vaccination. PBMCs from DV037 (Figure 3A) and BB232 (Figure 3B) were restimulated *in vitro* with the epitopic peptide 121-135. After CD8+ enrichment, cells were tested for cytotoxic activity against autologous B-LCLs sensitized with the epitopic peptide 121-135 (solid circles) or an irrelevant peptide (open circles). Figure 3C shows the results of an experiment where freshly isolated PBMCs from DV037, 51 weeks after its last immunization (two left panels), or *in vitro*-restimulated PBMCs from the same time point (two right panels) were restimulated for 12 hours with peptide 121-135 or a control peptide and stained for surface CD8 and intracellular IFN- $\gamma$  and TNF- $\alpha$ . Lymphocytes were gated by side vs. forward scatter light and then for CD8-PerCP. Plots show log fluorescence intensity for TNF- $\alpha$ -FITC and IFN- $\gamma$ -PE.

Figure 4 shows antibody titers against Core in the serum of immunized animals. Open bars, pre-immunization; striped bars, 2 weeks post 2<sup>nd</sup> immunization; filled bars, 2 weeks post 3<sup>rd</sup> immunization.

Figure 5 shows Th-1 and Th2-type cytokines in Core-ISCOM-immunized animals. The level of IFN- $\gamma$  (Figure 5A), IL-2 (Figure 5B), IL-5 (Figure 5C) and IL-10 (Figure 5D) was measured by specific ELISA in cell-free supernatant of freshly isolated

- 9 -

PBMCs stimulated for 48h as described in the examples. Open bars, pre-immunization; striped bars, 2 weeks post 2<sup>nd</sup> immunization; filled bars, 2 weeks post 3<sup>rd</sup> immunization. NT: Not Tested.

**Figure 6** shows MHC class I restriction of peptides 121-135 and 86-100 CTLs.

5 (Figure 6A) shows the results of an experiment where peptide 86-100-specific CTL

line derived from animal 15864 was tested in a standard <sup>51</sup>Cr release assay for its ability to lyse peptide 86-100 -sensitized B-LCL target cells derived from animals DV036 (solid squares), 15864 (solid circles), 15860 (solid triangles) and 15861 (open circles).

Figure 6B shows the results from an experiment where peptide 121-135-specific  
10 CTL line derived from animal 15862 was tested in a standard <sup>51</sup>Cr release assay for its ability to lyse peptide 121-135-sensitized B-LCL target cells derived from animals DV037 (solid squares), BB232 (solid triangles), 15862 (solid inverted triangles), 15863 (solid circles), 15861 (open squares) and 15860 (open inverted triangles).

15 **Figure 7** shows a quantification of the CD8+ and CD4+ T cell responses in Core-ISCOM-immunized animals. Freshly isolated PBMCs were restimulated *ex vivo* with rVVC/E1 or VVwt-infected autologous B-LCLs (Figure 7A) or with the recombinant Core protein of an *E. coli* control (Figure 7B). Cells were then stained for surface CD8 or CD4, and intracellular IFN- $\gamma$  and TNF- $\alpha$  as described in the examples.

20 Lymphocytes were gated by side vs. forward scatter light and then for CD8-PerCP (Figure 7A) or CD4-APC (Figure 7B). In Figure 7A, the corrected percent of CD8+ T cells with detectable IFN- $\gamma$  and/or TNF- $\alpha$  was calculated as (% CD8+ T cells restimulated with rVVC/E1 that were IFN- $\gamma$  and/or TNF- $\alpha$ +) – (% CD8+ T cells restimulated with VVwt that were IFN- $\gamma$  and/or TNF- $\alpha$ +) . In Figure 7B, the corrected  
25 percent of CD4+ T cells with detectable IFN- $\gamma$  and/or TNF- $\alpha$  was calculated as (% CD4+ T cells restimulated with Core that were IFN- $\gamma$  and/or TNF- $\alpha$ +) – (% CD4+ T cells restimulated with the *E. coli* that were IFN- $\gamma$  and/or TNF- $\alpha$ +) . Open bars, pre-immunization; striped bars, 2 weeks post 2<sup>nd</sup> immunization; filled bars, 2 weeks post 3<sup>rd</sup> immunization.

30 **Figure 8** shows that Core-ISCOM can serve as an adjuvant for E1E2. Mice (10 animals per group) were immunized with 2g of soluble E1E2 alone, 2 g of soluble

- 10 -

E1E2 + 2g of Core-ISCOM, or 2 g of soluble E1E2 adjuvanted with MF59. Mice were bled two weeks post 3<sup>rd</sup> immunization. Anti-E2 (filled bars) and anti-CD81 titers (striped bars) are presented as the geometric mean of the titers obtained from the individual mice from each group. NT: Not Tested.

5

**Figure 9** is a diagrammatic representation of the HCV genome, depicting the various regions of the polyprotein from which the present proteins for use with the ISCOMs are derived.

- 10 **Figure 10** is a graphical representation of the sucrose gradient analysis of NS35 Core 121-ISCOM™ formulations for NS35 Core 121-ISCOM™. (Figure 10A) and NS35 Core 121 protein (Figure 10B).

#### DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

15

The present invention is predicated, in part, on the development of an immunogenic complex formulation which utilises electrostatic interactions to associate an antigen of HCV and a carrier thereby facilitating, *inter alia*, the co-delivery of these molecules to the immune system. The immunogenic complexes of the present invention are particularly suitable for use in facilitating the stimulation of cytotoxic T-lymphocyte responses.

Accordingly, one aspect of the present invention relates to an immunogenic complex comprising a charged organic carrier and a charged antigen, which organic carrier and antigen are electrostatically associated, and wherein the charged antigen is a polypeptide of Hepatitis C Virus (HCV), preferably the core protein of HCV, or a fragment thereof, or a fusion protein comprising said polypeptide or a fragment thereof.

- 30 Reference to a "complex" should be understood as describing an entity of two or more different interacting chemical components.

- 11 -

Reference to a "charged" organic carrier or antigen should be understood as a reference to an organic carrier or antigen which exhibits an overall positive electrical charge or an overall negative electrical charge. By "overall" is meant the summation of the individual positive and negative charges which a given molecule comprises.

- 5 Where the summation of the individual positive and negative charges results in overall electrical neutrality, the molecule is not regarded as "charged" within the context of the present invention. Preferably, the organic carrier comprises an overall negative charge.
- 

- 10 Accordingly, the present invention more particularly provides an immunogenic complex as described above, wherein the charged organic carrier is a negatively charged organic carrier.

- Reference to "electrostatically associated" is a reference to the organic carrier and  
15 the antigen being linked, bound or otherwise associated by means which include electrostatic interaction. Accordingly, it should be understood that in some instances the electrostatic interaction will be the only attractive force which results in complexing of the antigen and the organic carrier. However, in other instances the formation of the electrostatic interaction may also lead to, or be associated with, the  
20 formation of other interactive forces.

- The terms "polyprotein", "protein" and "polypeptide" are used interchangeably herein and refer to a polymer of amino acid residues and are not limited to a minimum length of the product. Thus, peptides, oligopeptides, dimers, multimers, and the like,  
25 are included within the definition. Both full-length proteins and fragments thereof are encompassed by the definition. The terms also include postexpression modifications of the polypeptide, for example, glycosylation, acetylation, phosphorylation and the like. Furthermore, for purposes of the present invention, a "polypeptide" refers to a protein which includes modifications, such as deletions, additions and substitutions  
30 (generally conservative in nature), to the native sequence, so long as the protein maintains the desired activity. These modifications may be deliberate, as through site-directed mutagenesis, or may be accidental, such as through mutations of hosts

- 12 -

which produce the proteins or errors due to PCR amplification.

In particular, as shown in Figure 9, several proteins are encoded by the HCV genome. The order and nomenclature of the cleavage products of the HCV polyprotein is as follows: NH<sub>2</sub>-C-E1-E2-NS2-NS3-NS4a-NS4b-NS5a-NS5b-COOH.

Initial cleavage of the polyprotein is catalyzed by host proteases which liberate three structural proteins, the N-terminal nucleocapsid protein (termed "Core") and two envelope glycoproteins, "E1" (also known as E) and "E2" (also known as E2/NS1), as well as nonstructural (NS) proteins that contain the viral enzymes. The NS regions are termed NS2, NS3, NS4 and NS5. NS2 is an integral membrane protein with proteolytic activity. NS2, either alone or in combination with NS3, cleaves the NS2-NS3 sissle bond which in turn generates the NS3 N-terminus and releases a large polyprotein that includes both serine protease and RNA helicase activities. The NS3 protease serves to process the remaining polyprotein. Completion of polyprotein maturation is initiated by autocatalytic cleavage at the NS3-NS4a junction, catalyzed by the NS3 serine protease. Subsequent NS3-mediated cleavages of the HCV polyprotein appear to involve recognition of polyprotein cleavage junctions by an NS3 molecule of another polypeptide. In these reactions, NS3 liberates an NS3 cofactor (NS4a), two proteins with unknown function (NS4b and NS5a), and an RNA-dependent RNA polymerase (NS5b).

As explained above, any of a number of HCV polypeptides derived from the HCV polyprotein may be used in the immunogenic complexes of the present invention. Thus, these complexes may contain polypeptides derived from the HCV Core nucleocapsid protein, a nonstructural protein, the E1 envelope protein, the E2 envelope protein, polypeptide fragments of any of such proteins, or combinations of such proteins. Such fragments may be polypeptides comprising epitopes recognizable by T cells. Preferred fragments comprise those fragments which are immunogenic when provided by themselves, or when included in the immunogenic complex of the present invention.

- 13 -

Preferably, the immunogenic complex of the present invention comprises the core protein of HCV, or an immunogenic fragment thereof. The core protein of HCV has a pI of 10, making it highly positively charged at neutral and acidic pH. Reference to the "core protein" of HCV should be understood as including a reference to derivatives and equivalents of the core protein.

The polypeptide for use in the immunogenic complex of the present invention need not be physically derived from HCV, but may be synthetically or recombinantly produced using conventional techniques of molecular biology, microbiology, recombinant DNA, and immunology, which are within the skill of the art. Such techniques are explained fully in the literature, e.g. Sambrook *Molecular Cloning : A Laboratory Manual, Second Edition* (1989); *DNA Cloning, Volumes I and II* (D.N. Glover, ed., 1985); *Oligonucleotide Synthesis* (M.J. Gait, ed., 1986); *Nucleic Acid Hybridization* (B.D. Hames & S.J. Higgins, eds., 1984); *Transcription and Translation* (B.D. Hames & S.J. Higgins, eds., 1984); *Animal Cell Culture* (R.I. Freshney, ed., 1986); *Immobilized Cells and Enzymes* (IRL Press, 1986); B. Perbal, *A Practical Guide to Molecular Cloning* (1984); the *Methods in Enzymology* series (Academic Press, Inc.), especially volumes 154 and 155; *Gene Transfer Vectors for Mammalian Cells* (J.H. Miller and M.P. Calos, eds., 1987, Cold Spring Harbor Laboratory); Mayer and Walker, eds., (1987), *Immunochemical Methods in Cell and Molecular Biology* (Academic Press, London); Scopes, (1987), *Protein Purification: Principles and Practice, Second Edition* (Springer-Verlag, N.Y.), and *Handbook of Experimental Immunology, Volumes I-IV* (D.M. Weir and C.C. Blackwell, eds., 1986).

Moreover, the polypeptide may be derived from any of the various known HCV strains, such as from strains 1, 2, 3 or 4 of HCV. A number of conserved and variable regions are known between these strains and, in general, the amino acid sequences of epitopes derived from these regions will have a high degree of sequence homology, e.g., amino acid sequence homology of more than 30%, preferably more than 40%, when the two sequences are aligned. Thus, for example, the term "Core" polypeptide refers to the native Core protein from any of the various HCV strains, as well as Core analogs, muteins and immunogenic fragments, as

- 14 -

defined further below.

Reference to "derivative and equivalents" should be understood as a reference to chemical equivalents, mutants, homologs and analogs from natural, synthetic or  
5 recombinant sources. Derivatives may be derived from insertion, deletion or  
~~substitution of amino acids. Amino acid insertional derivatives include amino and/or~~  
carboxylic terminal fusions as well as intrasequence insertions of single or multiple  
amino acids. Insertional amino acid sequence variants are those in which one or  
more amino acid residues are introduced into a predetermined site in the protein  
10 although random insertion is also possible with suitable screening of the resulting  
product. Deletional variants are characterised by the removal of one or more amino  
acids from the sequence. Substitutional amino acid variants are those in which one  
residue in the sequence has been removed and a different residue inserted in its  
place. "Equivalents" can act as a functional analog of the subject antigen. Chemical  
15 equivalents may not necessarily be derived from the subject antigen but may share  
certain conformational similarities. Alternatively, chemical equivalents may be  
designed to mimic certain physiochemical properties of the subject antigen.  
Equivalents may be chemically synthesised or may be detected following, for  
example, natural product screening. Homologs contemplated herein include, but are  
20 not limited to, molecules derived from different species.

The present invention also extends to an immunogenic complex as described above  
wherein the charged antigen is a fragment of an HCV protein. By "fragment" is  
intended a polypeptide consisting of only a part of the intact full-length protein  
25 sequence and structure. The fragment can include a C-terminal deletion and/or an  
N-terminal deletion of the native polypeptide. An "immunogenic fragment" of a  
particular HCV protein will generally include at least about 5-10 contiguous amino  
acid residues of the full-length molecule, preferably at least about 15-25 contiguous  
amino acid residues of the full-length molecule, and most preferably at least about  
30 20-50 or more contiguous amino acid residues of the full-length molecule, that define  
an epitope, or any integer between 5 amino acids and the full-length sequence,  
provided that the fragment in question retains the ability to elicit an immune response

- 15 -

as defined below. For example, preferred immunogenic fragments, include but are not limited to fragments of the core protein of HCV that comprise, e.g., amino acids 10-45, 10-53, 67-88, 81-130, 86-100, 120-130, 121-135 and 121-170 of the polyprotein, numbered relative to the HCV-1a sequence presented in Choo *et al.* 5 (1991) *Proc Natl Acad Sci USA* 88:2451, as well as defined epitopes derived from the c33c region of the HCV polyprotein, as well as any of the other various epitopes identified from the HCV core, E1, E2, NS3 and NS4 regions. See, e.g., Chien *et al.* *Proc. Natl. Acad. Sci. USA* (1992) 89:10011-10015; Chien *et al.* *J. Gastroent. Hepatol.* (1993) 8:S33-39; Chien *et al.* International Publ. No. WO 93/00365; Chien, 10 D.Y. International Publ. No. WO 94/01778; allowed U.S. Patent Application Serial Nos. 08/403,590 and 08/444,818.

The term "epitope" as used herein refers to a sequence of at least about 3 to 5, preferably about 5 to 10 or 15, and not more than about 1,000 amino acids (or any 15 integer therebetween), which define a sequence that by itself or as part of a larger sequence, will stimulate a host's immune system to make a cellular antigen-specific immune response when the antigen is presented, or a humoral antibody response.

An epitope for use in the subject invention is not limited to a polypeptide having the exact sequence of the portion of the parent protein from which it is derived. Indeed, 20 viral genomes are in a state of constant flux and contain several variable domains which exhibit relatively high degrees of variability between isolates. Thus the term "epitope" encompasses sequences identical to the native sequence, as well as modifications to the native sequence, such as deletions, additions and substitutions (generally conservative in nature).

25 Regions of a given polypeptide that include an epitope can be identified using any number of epitope mapping techniques, well known in the art. See, e.g., *Epitope Mapping Protocols* in *Methods in Molecular Biology*, Vol. 66 (Glenn E. Morris, Ed., 1996) Humana Press, Totowa, New Jersey. For example, linear epitopes may be 30 determined by e.g., concurrently synthesizing large numbers of peptides on solid supports, the peptides corresponding to portions of the protein molecule, and reacting the peptides with antibodies while the peptides are still attached to the



- 16 -

- supports. Such techniques are known in the art and described in, e.g., U.S. Patent No. 4,708,871; Geysen *et al.* (1984) *Proc. Natl. Acad. Sci. USA* 81:3998-4002; Geysen *et al.* (1986) *Molec. Immunol.* 23:709-715, all incorporated herein by reference in their entireties. Similarly, conformational epitopes are readily identified
- 5 by determining spatial conformation of amino acids such as by, e.g., x-ray crystallography and 2-dimensional nuclear magnetic resonance. See, e.g., *Epitope Mapping Protocols, supra*. Antigenic regions of proteins can also be identified using standard antigenicity and hydropathy plots, such as those calculated using, e.g., the Omega version 1.0 software program available from the Oxford Molecular Group.
- 10 This computer program employs the Hopp/Woods method, Hopp *et al.*, *Proc. Natl. Acad. Sci USA* (1981) 78:3824-3828 for determining antigenicity profiles, and the Kyte-Doolittle technique, Kyte *et al.*, *J. Mol. Biol.* (1982) 157:105-132 for hydropathy plots.
- 15 As used herein, the term "conformational epitope" refers to a portion of a full-length protein, or an analog or mutein thereof, having structural features native to the amino acid sequence encoding the epitope within the full-length natural protein. Native structural features include, but are not limited to, glycosylation and three dimensional structure. Preferably, a conformational epitope is produced recombinantly and is
- 20 expressed in a cell from which it is extractable under conditions which preserve its desired structural features, e.g. without denaturation of the epitope. Such cells include bacteria, yeast, insect, and mammalian cells. Expression and isolation of recombinant conformational epitopes from the HCV polyprotein are described in e.g., International Publication Nos. WO 96/04301, WO 94/01778, WO 95/33053, WO
- 25 92/08734, which applications are herein incorporated by reference in their entirety.

As used herein the term "T-cell epitope" refers to a feature of a peptide structure which is capable of inducing T-cell immunity towards the peptide structure or an associated hapten. T-cell epitopes generally comprise linear peptide determinants

30 that assume extended conformations within the peptide-binding cleft of MHC molecules, (Unanue *et al.*, *Science* (1987) 236:551-557). Conversion of polypeptides

- 17 -

- to MHC class II-associated linear peptide determinants (generally between 5 - 14 amino acids in length) is termed "antigen processing" which is carried out by antigen presenting cells (APCs). More particularly, a T-cell epitope is defined by local features of a short peptide structure, such as primary amino acid sequence properties involving charge and hydrophobicity, and certain types of secondary structure, such as helicity, that do not depend on the folding of the entire polypeptide. Further, it is believed that short peptides capable of recognition by helper T-cells are generally amphipathic structures comprising a hydrophobic side (for interaction with the MHC molecule) and a hydrophilic side (for interacting with the T-cell receptor), (Margalit et al., *Computer Prediction of T-cell Epitopes*, New Generation Vaccines Marcel-Dekker, Inc, ed. G.C. Woodrow et al., (1990) pp. 109-116) and further that the amphipathic structures have an  $\alpha$ -helical configuration (see, e.g., Spouge et al. *J. Immunol.* (1987) 138:204-212; Berkower et al. *J. Immunol.* (1986) 136:2498-2503).
- Hence, segments of proteins which include T-cell epitopes can be readily predicted using numerous computer programs. (See e.g., Margalit et al., *Computer Prediction of T-cell Epitopes*, New Generation Vaccines Marcel-Dekker, Inc, ed. G.C. Woodrow et al., (1990) pp. 109-116). Such programs generally compare the amino acid sequence of a peptide to sequences known to induce a T-cell response, and search for patterns of amino acids which are believed to be required for a T-cell epitope.

- In general, "identity" refers to an exact nucleotide-to-nucleotide or amino acid-to-amino acid correspondence of two polynucleotides or polypeptide sequences, respectively. Percent identity can be determined by a direct comparison of the sequence information between two molecules by aligning the sequences, counting the exact number of matches between the two aligned sequences, dividing by the length of the shorter sequence, and multiplying the result by 100. Readily available computer programs can be used to aid in the analysis, such as ALIGN, Dayhoff, M.O. in *Atlas of Protein Sequence and Structure* M.O. Dayhoff ed., 5 Suppl. 3:353-358, National biomedical Research Foundation, Washington, DC, which adapts the local homology algorithm of Smith and Waterman *Advances in Appl. Math.* 2:482-489,

- 18 -

1981 for peptide analysis. Programs for determining nucleotide sequence identity are available in the Wisconsin Sequence Analysis Package, Version 8 (available from Genetics Computer Group, Madison, WI) for example, the BESTFIT, FASTA and GAP programs, which also rely on the Smith and Waterman algorithm. These  
5 programs are readily utilized with the default parameters recommended by the manufacturer and described in the Wisconsin Sequence Analysis Package referred to above. For example, percent identity of a particular nucleotide sequence to a reference sequence can be determined using the homology algorithm of Smith and Waterman with a default scoring table and a gap penalty of six nucleotide positions.

10

Another method of establishing percent identity in the context of the present invention is to use the MPSRCH package of programs copyrighted by the University of Edinburgh, developed by John F. Collins and Shane S. Sturrok, and distributed by IntelliGenetics, Inc. (Mountain View, CA). From this suite of packages the Smith-  
15 Waterman algorithm can be employed where default parameters are used for the scoring table (for example, gap open penalty of 12, gap extension penalty of one, and a gap of six). From the data generated the "Match" value reflects "sequence identity."

Other suitable programs for calculating the percent identity or similarity between sequences are generally known in the art, for example, another alignment program  
20 is BLAST, used with default parameters. For example, BLASTN and BLASTP can be used using the following default parameters: genetic code = standard; filter = none; strand = both; cutoff = 60; expect = 10; Matrix = BLOSUM62; Descriptions = 50 sequences; sort by = HIGH SCORE; Databases = non-redundant, GenBank + EMBL + DDBJ + PDB + GenBank CDS translations + Swiss protein + Spupdate +  
25 PIR. Details of these programs can be found at the following internet address: <http://www.ncbi.nlm.gov/cgi-bin/BLAST>.

The sequences for these polypeptides and proteins are known (see, e.g., U.S. Patent No. 5,350,671, incorporated herein by reference in its entirety). For example, a  
30 number of general and specific immunogenic polypeptides, derived from the HCV polyprotein, have been described. See, e.g., Houghton et al., European Publ. Nos. 318,216 and 388,232; Choo et al. *Science* (1989) 244:359-362; Kuo et al. *Science*

- 19 -

(1989) 244:362-364; Houghton et al. *Hepatology* (1991) 14:381-388; Chien et al. *Proc. Natl. Acad. Sci. USA* (1992) 89:10011-10015; Chien et al. *J. Gastroent. Hepatol.* (1993) 8:S33-39; Chien et al., International Publ. No. WO 93/00365; Chien, D.Y., International Publ. No. WO 94/01778. These publications provide an extensive  
5 background on HCV generally, as well as on the manufacture and uses of HCV polypeptide immunological reagents. For brevity, therefore, the disclosure of these  
publications is incorporated herein by reference.

It should be noted that for convenience, the various HCV regions are generally  
10 defined with respect to the amino acid number relative to the polypeptide encoded by the genome of HCV-1a, as described in Choo et al. (1991) *Proc Natl Acad Sci USA* 88:2451, with the initiator methionine being designated position 1. However, the polypeptides for use with the present invention are not limited to those derived from the HCV-1a sequence. In this regard, the corresponding regions in another HCV  
15 isolate can be readily determined by aligning sequences from the two isolates in a manner that brings the sequences into maximum alignment.

For example, HCV polypeptides derived from the Core region, such as polypeptides derived from the region found between amino acids 1-191; amino acids 10-53; amino  
20 acids 10-45; amino acids 67-88; amino acids 86-100; 81-130; amino acids 121-135; amino acids 120-130; amino acids 121-170; and any of the Core epitopes identified in, e.g., Houghton et al., U.S. Patent No. 5,350,671; Chien et al. *Proc. Natl. Acad. Sci. USA* (1992) 89:10011-10015; Chien et al. *J. Gastroent. Hepatol.* (1993) 8:S33-39; Chien et al., International Publ. No. WO 93/00365; Chien, D.Y., International  
25 Publ. No. WO 94/01778; and allowed U.S. Patent Application Serial Nos. 08/403,590 and 08/444,818, the disclosures of which are incorporated herein by reference in their entireties, will find use in the immunogenic complexes of the present invention.

HCV polypeptides derived from the envelope of HCV, including polypeptides derived  
30 from the E1 and E2 regions, as well as fusions between E1 and E2 will also find use herein. Particularly, the HCV envelope glycoproteins E1 and E2 form a stable

- 20 -

complex that is co-immunoprecipitable (Grakoui *et al.* (1993) *J. Virol.* **67**:1385-1395; Lanford *et al.* (1993) *Virology* **197**:225-235; Ralston *et al.* (1993) *J. Virol.* **67**:6753-6761). The HCV E1 and E2 glycoproteins have been shown to be protective in primate studies. (Choo *et al.* (1994) *Proc. Natl. Acad. Sci. USA* **91**:1294-1298).

5 The mature E1 region of HCV1 begins at approximately amino acid 192 of the polypeptide and continues to approximately amino acid 383. The mature E2 region of HCV1 begins at approximately amino acid 384-385 and extends as far as approximately amino acid residue 746 (see, Lin *et al.* *J. Virol.* (1994) **68**:5063-5073).

10 Additionally, polypeptides derived from the nonstructural regions of the virus will also find use herein. The NS3/4a region of the HCV polypeptide has been described and the amino acid sequence and overall structure of the protein are disclosed in Yao *et al.* *Structure* (November 1999) **7**:1353-1363. See, also, Dasmahapatra *et al.*, U.S. Patent No. 5,843,752, incorporated herein by reference in its entirety. As explained  
15 above, either the native sequence or immunogenic analogs can be used in the subject formulations. Dasmahapatra *et al.*, U.S. Patent No. 5,843,752 and Zhang *et al.*, U.S. Patent No. 5,990,276, both describe analogs of NS3/4a and methods of making the same.

20 Additionally, multiple epitope fusion antigens (termed "MEFAs"), as described in International Publ. No. WO 97/44469, may be associated with the immunogenic complexes. Such MEFAs include multiple epitopes derived from two or more of the various viral regions. The epitopes are preferably from more than one HCV strain, thus providing the added ability to protect against multiple strains of HCV in a single  
25 vaccine.

Moreover, polypeptides for use in the immunogenic complexes of this invention may be derived from the NS3 region of the HCV polypeptide. A number of such polypeptides are known, including, but not limited to polypeptides derived from the  
30 c33c and c100 regions, as well as fusion proteins comprising an NS3 epitope, such as c25. These and other NS3 polypeptides are useful in the present compositions and are known in the art and described in, e.g., Houghton *et al.*, U.S. Patent No.

- 21 -

5,350,671; Chien *et al. Proc. Natl. Acad. Sci. USA* (1992) 89:10011-10015; Chien *et al. J. Gastroent. Hepatol.* (1993) 8:S33-39; Chien *et al.*, International Publ. No. WO 93/00365; Chien, D.Y., International Publ. No. WO 94/01778; and allowed U.S. Patent Application Serial Nos. 08/403,590 and 08/444,818, the disclosures of which  
5 are incorporated herein by reference in their entireties.

---

It is readily apparent that a multitude of HCV polypeptides may be used in the immunogenic complex formulations or may be coadministered therewith, in order to provide a cellular immune response against the HCV antigen in question.

10

The present invention further extends to an immunogenic complex as described above wherein the charged protein is a fusion protein comprising the core or another protein of HCV or a fragment thereof. Reference to a "fusion protein" should be understood as a reference to a fusion in which the HCV core or other protein or  
15 fragment is operatively linked to another peptide, polypeptide or protein. The term "operatively linked" is intended to indicate that the HCV core or other protein or fragment and the other peptide, polypeptide or protein are fused in-frame to each other, either directly or indirectly through a linker peptide or polypeptide, with the fusion being at either the N-terminal end or the C-terminal end of the HCV core or  
20 other protein or fragment.

The fusion protein may comprise a tag protein or peptide moiety such as a hexa-his (His)<sub>6</sub> moiety, glutathione-S-transferase (GST) moiety or a FLAG moiety.

25 Preferably, however, the fusion protein comprises a second immunogenically active peptide, polypeptide or protein which may be derived from HCV, or some other viral, bacterial, fungal or similar organism.

The linker peptide or polypeptide, where present in the fusion protein, may comprise  
30 a sequence of from 1 to 50, preferably 1 to 20, and more preferably 1 to 5 amino acid residues.

- 22 -

- Reference throughout this specification to "organic carrier" should be understood as a reference to any molecule, aggregate or complex of molecules, compound or other entity which, when an antigen is associated with it, facilitates the induction of an immune response, and in particular a cytotoxic T-lymphocyte response, to the antigen. The subject carrier is "organic" and, in this regard, "organic" should be understood as a compound of carbon whether naturally, recombinantly or synthetically obtained or derived. In a particularly preferred embodiment the organic carrier is an adjuvant. By "adjuvant" is meant any molecule, aggregate or complex of molecules, compound or other entity which functions to stimulate, enhance or otherwise up-regulate any one or more aspects of the immune response. For example, the adjuvant may induce inflammation thereby attracting immune response cells to the site of antigen localisation. Alternatively, the adjuvant may slowly release the antigen thereby providing on-going stimulation of the immune system.
- Examples of charged organic carriers which are adjuvants suitable for use in the present invention include, but are not limited to, saponin, saponin complexes, any one or more components of the immunostimulating complex of saponin, cholesterol and lipid known as ISCOMATRIX™ (for example the saponin component and/or the phospholipid component), liposomes or oil-in-water emulsions. [The composition and preparation of ISCOMATRIX™ is described in detail in International Patent Application Number PCT/SE86/00480, Australian Patent Numbers 558258 and 632067 and European Patent Publication No. 0 180 564, the disclosures of which are incorporated herein by reference]. Further examples of adjuvants include, but are not limited to, those detailed in the publications of Cox and Coulter, 1992, 1997 and 1999. It should be understood that the subject organic carrier may be naturally occurring or it may be synthetically or recombinantly derived.

Accordingly, the present invention still more preferably provides an immunogenic complex as described above, wherein the charged organic carrier is a negatively charged adjuvant.

- 23 -

Preferably, said adjuvant comprises saponin or a saponin complex. More preferably, said saponin complex is ISCOMATRIX™.

The organic carrier of the present invention may also be, in its initial or natural form, negatively charged, positively charged or neutral. Increasing the degree of negative charge (for example, where the organic carrier is only weakly negatively charged) or converting a neutral or positively charged organic carrier to a negatively charged organic carrier may also be achieved by any suitable method known to those skilled in the art. For example, where the organic carrier is an oil-in-water emulsion, incorporation of any anionic surfactant with a non-polar tail will impart an overall negative charge to the emulsion due to insertion of the tail of the surfactant into the oil droplet which thereby leaves the negatively charged head group exposed. The negative charge of a saponin complex adjuvant may be increased, for example, by the addition of negatively charged lipid during complex formation.

Examples of detergents which can increase the negative charge of a carrier include, but are not limited to cholic acid, deoxycholic acid, taurocholic acid and taurodeoxycholic acid. Examples of lipids which can increase the negative charge of a carrier include, but are not limited to, phospholipids (preferably phosphatidyl inositol, phosphatidyl serine, phosphatidyl glycerol and phosphatidic acid and most preferably cardiolipin) and bacterial lipids (preferably monophosphoryl lipid A(MPL) and most preferably diphosphoryl lipid A, such as OM174 as described in International Patent Publication No. WO 95/14026).

Without limiting the present invention in any way, the inventors have determined that where the subject charged organic carrier and charged antigen are naturally negatively and positively charged, respectively, the object of the invention can be achieved. However, a still more effective immunogenic complex may be achieved if the subject naturally negatively charged organic carrier is rendered more negatively charged (preferably by addition of cardiolipin or diphosphoryl lipid A).



- 24 -

Accordingly, in one preferred embodiment there is provided an immunogenic complex as described above, wherein the negatively charged adjuvant is a naturally negatively charged adjuvant which has been modified to increase the degree of its negative charge.

5

~~Reference to an adjuvant being "naturally" negatively charged, should be understood~~  
as a reference to the charge which the molecule bears upon its creation - whether that be by natural, recombinant or synthetic means. Modification to increase the degree of charge can be achieved by any suitable technique as hereinbefore  
10 discussed. Preferably, the subject adjuvant is rendered more negative via the addition of cardiolipin or diphosphoryl lipid A.

The present invention is predicated, in part, on the formation of immunogenic complexes via the electrostatic association, preferably, of a negatively charged  
15 organic carrier with a positively charged antigen. The administration of such a complex to a subject facilitates the induction of a significantly better immune response than would be achieved were the adjuvant and antigen administered simultaneously but in a non-associated form. In particular, the administration of an antigen associated with an adjuvant, according to the present invention, facilitates the  
20 induction of a cytotoxic T-lymphocyte response to the antigen. However, humoral and other cellular responses can also be enhanced.

Generally, in the immunogenic complex of the present invention, the ratio of the charged organic carrier to the charged antigen, by weight, is in the range of 5:1 to  
25 0.5:1. Preferably, the ratio by weight is approximately 3:1 to 1:1, and more preferably the ratio by weight is 2:1.

Without limiting the present invention to any one theory or mode of action, it is thought that the complexing of the adjuvant with the antigen facilitates co-delivery of  
30 the adjuvant and the antigen to the same antigen presenting cell thereby facilitating the induction of immune responses which either would not occur or would not occur as effectively were these molecules not co-delivered. For example, the induction of

- 25 -

some CD8+ cytotoxic T-lymphocyte responses are thought to occur where the adjuvant induces endosomal escape of the antigen in the antigen presenting cell. This necessarily requires co-delivery of the antigen and the adjuvant to the antigen presenting cell.

5

A further aspect of the present invention therefore relates to the use of the invention to induce an immune response in a mammal including, but not limited to, a humoral and/or cell mediated immune response.

- 10 Accordingly, another aspect of the present invention relates to a vaccine composition comprising as the active component an immunogenic complex comprising a charged organic carrier and a charged antigen, which organic carrier and antigen are electrostatically associated, and wherein the charged antigen is a polyprotein of Hepatitis C Virus (HCV) or a fragment thereof, or a fusion protein comprising said
- 15 polyprotein or a fragment thereof, together with one or more pharmaceutically acceptable carriers and/or diluents.

Preferably, said organic carrier is an adjuvant, and even more preferably a saponin or a saponin complex. Preferably said saponin complex is ISCOMATRIX™.

20

Preferably said organic carrier is negatively charged.

- Vaccine compositions according to this aspect of the invention may be either prophylactic (i.e. used to prevent infection) or therapeutic (i.e. used to treat disease
- 25 after infection). The vaccine compositions are conventionally administered parenterally, e.g. by injection, either subcutaneously, intramuscularly, or transdermally/transcutaneously. Additional formulations suitable for other modes of administration include oral and pulmonary formulations, suppositories, and transdermal applications. Dosage treatment may be a single dose schedule or a
- 30 multiple dose schedule. The vaccine compositions may be administered in conjunction with other immunoregulatory agents.

- 26 -

These vaccine compositions comprise an immunogenic complex of the present invention in combination with one or more pharmaceutically acceptable carriers and/or diluents, such carriers include any carrier that does not itself induce the production of a response harmful to the individual receiving the composition. Suitable carriers are typically large, slowly metabolised macromolecules such as proteins, polysaccharides, polylactic acids, polyglycolic acids, polymeric amino acids, amino acid copolymers, lipid aggregates (such as oil droplets or liposomes), and inactive virus particles. Such carriers are well known to those of ordinary skill in the art. Additionally, these carriers may function as immunostimulating agents or adjuvants in addition to the adjuvant effect of the immunogenic complex itself. Furthermore, the antigen may be conjugated to a bacterial toxoid, such as a toxoid from diphtheria, tetanus, cholera, *H. pylori*, etc., pathogens.

The vaccine compositions may also include further adjuvants to enhance effectiveness of the composition. Suitable adjuvants include, but are not limited to: (1) aluminum salts (alum), such as aluminum hydroxide, aluminum phosphate, aluminum sulfate, etc; (2) oil-in-water emulsion formulations (with or without other specific immunostimulating agents such as muramyl peptides (see below) or bacterial cell wall components), such as for example (a) MF59 (PCT Publ. No. WO 90/14837), containing 5% Squalene, 0.5% Tween 80, and 0.5% Span 85 (optionally containing various amounts of MTP-PE (see below), although not required) formulated into submicron particles, (b) SAF, containing 10% Squalene, 0.4% Tween 80, 5% pluronic-blocked polymer, and thr-MDP (see below) either microfluidised into a submicron emulsion or vortexed to generate a large particle size emulsion, and (c) Ribi™ adjuvant system (RAS), (Ribi Immunochem, Hamilton, MT) containing 2% Squalene, 0.2% Tween 80, and one or more bacterial cell wall components from the group consisting of monophosphorylipid A (MPL), trehalose dimycolate (TDM), and cell wall skeleton (CWS), preferably MPL+CWS (Detox™); (3) saponin adjuvants, such as Stimulon™ (Cambridge Bioscience, Worcester, MA); (4) Complete Freund's Adjuvant (CFA) and Incomplete Freund's Adjuvant (IFA); (5) cytokines, such as interleukins (e.g. IL-1, IL-2, IL-4, IL-5, IL-6, IL-7, IL-12, etc), interferons (e.g. gamma interferon), macrophage colony stimulating factor (M-CSF), tumor necrosis factor

- 27 -

(TNF), etc.; (6) detoxified mutants of a bacterial ADP-ribosylating toxin such as a cholera toxin (CT), a pertussis toxin (PT), or an *E. coli* heat-labile toxin (LT), particularly LT-K63, LT-R72, CT-S109, PT-K9/G129; see, e.g. WO 93/13302 and WO 92/19265; (7) other substances that act as immunostimulating agents to  
5 enhance the effectiveness of the composition; and (8) microparticles with adsorbed macromolecules, as described in International Patent Application No. PCT/US99/17308. Alum and MF59 are preferred.

As mentioned above, suitable muramyl peptides include, but are not limited to, N-acetyl-muramyl-L-threonyl-D-isoglutamine (thr-MDP), N-acetyl-normauramyl-L-alanyl-D-isoglutamine (nor-MDP), N-acetylmuramyl-L-alanyl-D-isoglutaminyl-L-alanine-2-(1'-2'-dipalmitoyl-sn-glycero-3-hydroxyphosphoryloxy)-ethylamine (MTP-PE), etc.

The vaccine compositions typically will also contain diluents, such as water, saline,  
15 glycerol, ethanol, etc. Additionally, auxiliary substances, such as wetting or emulsifying agents, pH buffering substances, and the like, may be present in the compositions.

The forms suitable for injectable use include sterile aqueous solutions (where water  
20 soluble) or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersions. They must be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms such as bacteria and fungi. The prevention of the action of microorganisms can be brought about by various antibacterial and antifungal agents,  
25 for example, parabens, chlorobutanol, phenol, sorbic acid, thimerosal and the like.

In many cases, it will be preferable to include isotonic agents, for example, sugars or sodium chloride. Prolonged absorption of the injectable compositions can be brought about by the use in the compositions of agents delaying absorption, for example, aluminium monostearate and gelatin.

30

Sterile injectable solutions are prepared by incorporating the active compounds in the required amount in the appropriate solvent with various of the other ingredients

- 28 -

enumerated above, as required, followed by filter sterilisation. Generally, dispersions are prepared by incorporating the various sterilised active ingredients into a sterile vehicle which contains the basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and the freeze-drying technique which yield a powder of the active ingredient plus any additional desired ingredient from previously sterile-filtered solution thereof.

- 10 When the active ingredients are suitably protected they may be orally administered, for example, with an inert diluent or with an assimilable edible carrier, or they may be enclosed in hard or soft shell gelatin capsule, compressed into tablets, or incorporated directly with the food of the diet. For oral therapeutic administration, the active compound may be incorporated with excipients and used in the form of
- 15 ingestible tablets, buccal tablets, troches, capsules, elixirs, suspensions, syrups, wafers, and the like. Such compositions and preparations should contain at least 1% by weight of active compound. The percentage of the compositions and preparations may, of course, be varied and may conveniently be between about 5 to about 80% of the weight of the unit. The amount of active compound in such therapeutically
- 20 useful compositions is such that a suitable dosage will be obtained. Preferred compositions or preparations according to the present invention are prepared so that an oral dosage unit form contains between about 0.1  $\mu$ g and 2000 mg of active compound.
- 25 The tablets, troches, pills, capsules and the like may also contain the components as listed hereafter: a binder such as gum, acacia, corn starch or gelatin; excipients such as dicalcium phosphate; a disintegrating agent such as corn starch, potato starch, alginic acid and the like; a lubricant such as magnesium stearate; and a sweetening agent such as sucrose, lactose or saccharin may be added or a flavouring agent such
- 30 as peppermint, oil of wintergreen, or cherry flavouring. When the dosage unit form is a capsule, it may contain, in addition to materials of the above type, a liquid carrier. Various other materials may be present as coatings or to otherwise modify the

- 29 -

physical form of the dosage unit. For instance, tablets, pills, or capsules may be coated with shellac, sugar or both. A syrup or elixir may contain the active compound, sucrose as a sweetening agent, methyl and propylparabens as preservatives, a dye and flavouring such as cherry or orange flavour. Of course, any  
5 material used in preparing any dosage unit form should be pharmaceutically pure and substantially non-toxic in the amounts employed. In addition, the active compound(s) may be incorporated into sustained-release preparations and formulations.

---

Without limiting the operation of the present invention in any way, the co-delivery of  
10 the immunogenic complex of the present invention is particularly useful for inducing an immune response and, in particular, a cytotoxic T-lymphocyte response to an antigen. Said immune response may be a specific (T cell and/or B cell) and/or non-specific immune response.

15 Accordingly, still another aspect of the present invention relates to a method of eliciting, inducing or otherwise facilitating, in a mammal, an immune response to an antigen, said method comprising administering to said mammal an effective amount of an immunogenic complex or a vaccine composition as hereinbefore described.

20 Preferably said immune response comprises a cytotoxic T-lymphocyte response.

It should be understood that the subject cytotoxic lymphocyte response may occur either in isolation or together with a helper T cell response, a humoral response or other specific or non-specific immune response.

25

A further aspect of the present invention relates to the use of the immunogenic complex of the invention in relation to the therapeutic and/or prophylactic treatment of disease conditions. Examples of disease conditions which can be treated in accordance with the method of the present invention include any disease condition  
30 which results from HCV infection.

- 30 -

Accordingly, yet another aspect of the present invention relates to a method of treating a disease condition in a mammal, said method comprising administering to said mammal an effective amount of an immunogenic complex or a vaccine composition as hereinbefore described, wherein administering said composition  
5 elicits, induces or otherwise facilitates an immune response which inhibits, halts, ~~delays or prevents the onset or progression of the disease condition.~~

---

Direct delivery of the compositions will generally be accomplished by injection, either subcutaneously, intraperitoneally, intravenously or intramuscularly or delivered to the  
10 interstitial space of a tissue. The compositions can also be administered into a lesion. Other modes of administration include oral and pulmonary administration, suppositories, and transdermal or transcutaneous applications. Dosage treatment may be a single dose schedule or a multiple dose schedule.

15 An "effective amount" means an amount necessary at least partly to attain the desired immune response, or to delay the onset or inhibit progression or halt altogether, the onset or progression of a particular condition being treated. This amount varies depending upon the health and physical condition of the individual to be treated, the taxonomic group of individual to be treated, the capacity of the  
20 individual's immune system to synthesise antibodies, the degree of protection desired, the formulation of the vaccine, the assessment of the medical situation, and other relevant factors. It is expected that the amount will fall in a relatively broad range that can be determined through routine trials.

25 The term "mammal" includes humans, primates, livestock animals (eg. horses, cattle, sheep, pigs, donkeys), laboratory test animals (eg. mice, rats, rabbits, guinea pigs), companion animals (eg. dogs, cats) and captive wild animals (eg. kangaroos, deer, foxes). Preferably, the mammal is a human or laboratory test animal. Even more preferably, the mammal is a human.

- 31 -

The mammal undergoing treatment may be human or an animal in need of therapeutic or prophylactic treatment of a disease condition or a potential disease condition.

- 5 In yet another aspect the present invention relates to the use an immunogenic complex or vaccine composition as hereinbefore described in the manufacture of a medicament for inhibiting, halting, delaying or preventing the onset or progression of a disease condition.
- 
- 10 Yet another aspect of the present invention relates to an agent for use in inhibiting, halting, delaying or preventing the onset or progression of a disease condition. Said agent comprising an immunogenic complex or vaccine composition as hereinbefore described.
- 15 Further features of the present invention are more fully described in the following non-limiting Examples.

Reference to "ISCOPREP™ 703" should be understood as a reference to a saponin preparation comprising from 50-90% by weight of Fraction A of Quil A and 50% to  
20 10% by weight of Fraction C of Quil A. Fractions A and C are prepared from the lipophilic fraction of Quil A. Fractions "A" and "C", their method of preparation and the method of preparing ISCOPREP™ 703 are detailed in International Patent Publication No. WO96/11711, which is incorporated herein by reference.

- 25 Below are examples of specific embodiments for carrying out the present invention. The examples are offered for illustrative purposes only, and are not intended to limit the scope of the present invention in any way. Those of skill in the art will readily appreciate that the invention may be practiced in a variety of ways given the teaching of this disclosure.

30

Efforts have been made to ensure accuracy with respect to numbers used (e.g., amounts, temperatures, etc.), but some experimental error and deviation should, of



- 32 -

course, be allowed for.

### Materials and Methods

#### *Animals.*

- 5 Rhesus macaques (*Macaca mulatta*) were housed at Southwest Foundation for Biomedical Research (SFBR, San Antonio, TX). Studies were performed under the NIH Guidelines for Care and Use of Laboratory Animals (National Institute of Health. (1985) *Guide for the care and use of laboratory animals*. US department of Health and Human services. publication No 82-23. National Institute of Health, Bethesda, MD). Class I major histocompatibility complex (MHC) typing of the animals was performed as described (Urvater. *et al.* (2000) *J. Immunol.* **164**:1386.).
- 

- Female C57BL/6 (H-2<sup>b</sup>) mice were purchased from Charles River Laboratories and used between 8 and 10 weeks of age. Mice were housed in a pathogen free environment and were handled according to the international guidelines for experimentation with animals.

#### *Immunogens and Adjuvants.*

- The E. coli-derived full-length HCV-1a Core recombinant protein (aa: 1-191) was produced and purified under GMP conditions and is more than 98% pure. The recombinant HCV-1a E1E2<sub>809</sub> protein was produced in CHO cells. This modified E1E2 protein contained amino acids 192 to 809. The recombinant NS35Core121 protein was produced in yeast cells. The adjuvant LTK63 is a genetically detoxified mutant of the heat-labile enterotoxin of *Escherichia coli*, in which the Serine at position 63 is replaced by a Lysine (Partidos *et al.* (1999) *Immunol. Lett.* **67**:209.). The Core-ISCOM formulations were prepared by mixing the core protein with a preformed ISCOMATRIX™ (empty ISCOMs™) utilizing ionic interactions to maximize association between the antigen and the adjuvant. ISCOMATRIX™ was prepared essentially by previously described methods except that diafiltration was used in place of dialysis

- 33 -

(Coulter *et al.* (1998) *Vaccine* 16:1243). The oil-in-water adjuvant MF59 has been described (Ott *et al.* (1995) *Pharm. Biotechnol.* 6:277).

#### *Sucrose gradient analysis of Core-ISCOM*

- 5 After formulation, preparations were purified on a sucrose gradient (10 to 50% w/v) and fractions analysed for ISCOMATRIX™ and protein to determine the amount of association. ISCOMATRIX™ was analysed using diphenylhexatriene (DPH) which fluoresces when associated with lipid. Briefly, DPH was dissolved at 1mg/ml in acetone then diluted 1 in 50 in PBS pH7.2, then 50µl mixed with 50µl of each fraction in a
- 10 microtitre plate. Following incubation for 150 mins at 20-25°C the plate was read in a fluorometer using excitation 355nm and emission 460nm. Protein was detected using Pierce Coomassie according to manufacturers instructions. Briefly, 50µl Coomassie solution was added to 50µl of each fraction in a microtitre plate. The plate was mixed and absorbance read at 595nm.

15

#### *Particle size analysis*

Formulations were analysed for particle size by dynamic light scattering using a Nicomp Submicron Particle Sizer Model 370.

20

#### *Peptides and Vaccinia Viruses.*

- Peptides (15 or 20mer overlapping by 10 aa) spanning the entire length of the Core (aa: 1-191) protein of HCV-1a (Choo *et al.* (1991) *Proc Natl Acad Sci USA* 88:2451) were synthesized with free amine N-termini and free acid C-termini by Research Genetics (Huntsville, AL). The recombinant vaccinia virus (rVV) expressing Core and
- 25 E1 (aa: 1-384; rVVC/E1) and wild type VV (VVwt) have been described (Cooper *et al.* (1999) *Immunity* 10:439).

#### *Immunization.*

- Rhesus macaques were immunized under anesthesia. The first study was comprised
- 30 of nine animals divided into three groups of three animals each. The first group

- 34 -

(animals BB228, BB232 and DV036) were infected with  $2 \times 10^8$  plaque forming units (pfu) ( $1 \times 10^8$  intradermally and  $1 \times 10^8$  by scarification) of rVVC/E1 at month 0. This group served as a positive control for CTL priming. Animals from the second group (AY921, BB231 and DV037) were immunized with 25 $\mu$ g of Core-ISCOM by intramuscular (IM) injection in the left quadriceps at month 0, 1, 2 and 6. Animals from the third group (AY922, BB227 and BB230) were immunized by IM injection with 200 $\mu$ g of HCV-Core protein adjuvanted with 200 $\mu$ g of LTK63 at month 0, 1, 2 and 6. For the second study, five animals (15860, 15861, 15862, 15863 and 15864) were immunized with 50 $\mu$ g of Core-ISCOM by IM injection in the left quadriceps at month 0, 1 and 2. Some Core-immunized animals (see Table I) also received  $2 \times 10^8$  pfu ( $1 \times 10^8$  intradermally and  $1 \times 10^8$  by scarification) of rVVC/E1 nine or eleven weeks post their last vaccine immunization.

Mice (10 animals per group) were immunized in the tibialis anterior muscles (50 $\mu$ l per muscle) with 2 $\mu$ g per dose of recombinant E1E2 protein alone or 2 $\mu$ g per dose of recombinant E1E2 protein in the presence of MF59 (vol:vol), or 2 $\mu$ g per dose of recombinant E1E2 protein + 2 $\mu$ g per dose of Core-ISCOM at weeks 0, 4 and 8.

#### *Cells and cell lines.*

Peripheral blood was drawn from the femoral vein while the animals were under anesthesia. PBMCs were obtained after centrifugation over a Ficoll-hypaque gradient and cultured in 24-well dishes at  $5 \times 10^6$  cells per well. Of those cells,  $1 \times 10^6$  were sensitized with 10 $\mu$ M of a peptide pool (consisting of individual peptides) for one hour at 37°C, washed and added to the remaining  $4 \times 10^6$  untreated PBMCs in 2 ml of culture medium (RPMI 1640, 10% heat-inactivated fetal bovine serum, and 1% antibiotics) supplemented with 10 ng/ml of IL-7 (R&D, Minneapolis, MN). After 48 hours, 5% (final) of interleukin-2 containing supernatant (T-STIM without PHA, Collaborative Biomedical Products, Bedford, MA) and 50 U/ml (final) of recombinant IL-2 (Chiron Corporation, Emeryville, CA) were added to the cultures. Cultures were fed every 3-4 days. After 10 days in culture, CD8<sup>+</sup> T cells were isolated using anti-CD8 antibodies bound to magnetic beads (Dynal, Oslo, Norway) according to the manufacturer's instructions. Purified CD8<sup>+</sup> cells (>93% pure as determined by flow

- 35 -

cytometry) were cultured for another 2-3 days prior to being assayed for cytotoxic activity. Peptide-specific CD8<sup>+</sup> lines were obtained by periodically restimulating these CD8<sup>+</sup> T cells with autologous B-LCLs and peptide.

- 5 B-LCLs were derived from each animal using supernatants from the H. papio producer cell line S394.
- 

*CTL assay.*

Cytotoxic activity was assayed in a standard <sup>51</sup>Cr release assay as described  
10 elsewhere (Paliard *et al.* (2000) *AIDS Res. Hum. Retroviruses* 16:273). Briefly, B-LCLs were incubated with 10 $\mu$ M of peptides and 50 $\mu$ Ci of <sup>51</sup>Cr for 1h, washed three times and plated at 5 x 10<sup>3</sup> cells per well in a 96 well plate. Alternatively, B-LCLs were infected at multiplicity of infection (MOI) of 10:1 with rVVC/E1 or VVwt for 1h, washed and cultured overnight prior to labeling with <sup>51</sup>Cr. CD8<sup>+</sup> cells were plated in  
15 duplicate at three different E:T ratio and incubated with target cells for 4 hours in the presence of 2 x 10<sup>5</sup> per well of unlabeled target cells (cold targets), that were added to minimize lysis of B-LCLs by H. papio or endogenous virus (e.g. foamy virus)-specific CTLs. CTL responses were scored positive when percent specific lysis at the two highest E:T ratios were greater than or equal to the percent of lysis of control  
20 targets plus 10.

*Lymphoproliferation assay.*

This assay has been described previously (Hong *et al.* (1997) *J. Virol.* 71:6427). Briefly, freshly isolated PBMCs were plated in triplicates at 2 x 10<sup>5</sup> cells per well in 96  
25 well round bottomed plates and cultured in the presence of 5 $\mu$ g/ml of recombinant Core protein or 0.05  $\mu$ g/ml of *E. coli* control. Plates were pulsed with 1 $\mu$ Ci per well of <sup>3</sup>H-thymidine on day 5 and harvested 6-8 hours later. Results are presented as stimulation index (SI) calculated as (mean experimental cpm) / (mean cpm in the presence of the *E. coli* control). An SI  $\geq$ 3.0 was scored positive.

- 36 -

**FACS analysis.**

- Freshly isolated PBMCs or PBMCs that had been restimulated *in vitro* with a peptide were cultured in media alone or restimulated with 5 µg/ml of Core protein, 0.05 µg/ml of *E. coli* control, 5 µg/ml of peptide, or VV-infected or peptide-sensitized autologous
- 5 B-LCLs (1:1) for 12 hours in culture media containing 50 U/ml of rIL-2 (Chiron Corporation, Emeryville, CA) and 3 µM monensin (Pharmingen, San Diego, CA).

- Cells were stained according to Pharmingen's protocol for surface CD4 and CD8 with APC-conjugated anti-human CD4 and PerCP-conjugated anti-human CD8, and for intracellular IFN-γ and TNF-α with PE-conjugated anti-human IFN-γ and FITC-
- 10 conjugated anti-human TNF-α. Antibodies were from Pharmingen and Becton-Dickinson (San Jose, CA). Cells were analyzed on a FACScalibur. Data files were analyzed using the CellQuest software.

**Cytokine ELISA.**

- 15 Freshly isolated Rhesus macaque's PBMCs were restimulated with peptides encompassing the whole Core protein. Levels of Rhesus monkey IL-2, IL-5, IL-10 and IFN-γ present in 48 hours cell-free culture supernatants were determined by specific ELISA (U-Cytech, Utrecht, The Netherlands) following the manufacturers' specification.

20

**HCV antibodies.**

- Serum levels of HCV Core and HCV E2 antibodies were quantified by ELISA as described (Chien *et al.* (1992) *Proc Natl Acad Sci USA* 89:10011). Serum levels of antibodies inhibiting the binding of E2 to the putative HCV receptor CD81 (Pileri *et*
- 25 *al.* (1998) *Science* 282:938) were determined by immunoassay.

**Example 1****Sucrose Gradient Analysis of Core-ISCOM**

- Core Protein was found in fractions 5 to 11 (Figure 1A) whilst ISCOMATRIX® was
- 30 found in fractions 9 to 13 (Figure 1B). The Core-ISCOM was found with in fractions 14-

- 37 -

17 with both the ISCOM™ and the protein peaks overlapping, which indicates association (Figure 1C).

## Example 2

### Stability of Core-ISCOM

The stability of the Core-ISCOM™ formulation was evaluated for 11 months. Both the particle size and association remained consistent for at least 11 months at 2-8°C (Table I).

## Example 3

### Priming of Core-specific CTLs in Vaccinated Animals

As explained above, two different prototype vaccines (Core-ISCOM and Core + LTK63) aimed at eliciting HCV-Core-specific CTLs were each administered to three HCV-naïve Rhesus macaques (see Table II for animal assignment, dosage and immunization schedule). Since it was unknown whether Rhesus macaques' MHC class I molecules can bind and present HCV-Core -derived peptides and whether the positively selected CD8+ T cell repertoire in these animals can recognize such MHC class I – Core-derived peptide complexes, three additional animals were inoculated with  $2 \times 10^8$  pfu of rVVC/E1 to serve as positive controls (Table II).

None of the nine animals had any detectable CTLs at the time of immunization (week 0; Table III and data not shown). This confirmed that these animals had not been previously exposed to HCV Core and that restimulation of PBMCs under the conditions described in Material and Methods did not result in the priming of primary CTL responses *in vitro*. Two weeks post rVVC/E1 infection, two (BB232 and DV036) out of the three animals had detectable CTLs against Core peptides pool 4 (aa: 121-170) and pool 3 (aa: 81-130), respectively (Table III). By deconvoluting these peptide pools, it was determined that BB232's CTLs recognized the epitopic peptide 121-135 and that DV036's CTLs recognized peptide 86-100. The presence of 121-135 and

- 38 -

86-100-specific CTLs in these rVVC/E1-inoculated animals indicated that both peptides were naturally processed. No CTL responses were detectable in the other rVVC/E1-inoculated animal (BB228). This indicated that Core-specific CTLs can be elicited in at least some Rhesus monkeys. Out of the three animals that received Core adjuvanted with LTK63 (Table II), only one animal (BB230) showed a CTL response against Core. This response, directed against pool 2 (aa: 41-90), was transient, however, as it was detectable two weeks post 3<sup>rd</sup> immunization, but was undetectable six weeks post 3<sup>rd</sup> immunization. Furthermore, these CTLs were not boosted by a 4<sup>th</sup> immunization (Table III) and the individual peptide recognized could not be identified. Two out of the three Core-ISCOM-immunized animals (AY921 and BB231; Table II) did not mount a detectable Core-specific CTL response. In contrast, in the other Core-ISCOM-immunized animal (DV037), CTLs recognizing pool 4 (aa: 121-170) were detectable as early as two weeks post 2<sup>nd</sup> immunization. This response was directed against the epitopic peptide aa: 121-135 and was also present post 3<sup>rd</sup> and post 4<sup>th</sup> immunization (Table III).

In contrast to animals that received the Core + LTK63 prototype vaccine, animals immunized with the Core-ISCOM prototype vaccine had detectable CTLs post second immunization and these CTLs were also present post 3<sup>rd</sup> and 4<sup>th</sup> immunization (Table III). Thus, this latter vaccine seemed more potent at eliciting CTL responses in Rhesus macaques than the former one. However, this formulation primed Core-specific CTLs in only one out of the three animals. This might be due to the fact that the MHC class I molecules of the non-responding animals were unable to bind and present peptides derived from this relatively small protein (191 aa). To test this hypothesis, CTL lines specific for peptide 121-135 and 86-100 were established from responding animals. As shown in Figure 2A, the peptide 121-135-specific CTL line lysed peptide-sensitized B-LCLs derived from DV037, but did not kill peptide 121-135-sensitized B-LCLs from the two non-responding animals (AY921 and BB231). Similarly, B-LCLs derived from DV036 but not AY921 or BB231 were able to present peptide 86-100 to CD8+ CTLs (Figure 2B). These data indicated that AY921's and BB231's MHC class I molecules could not present these peptides to CD8+ T cells, and suggested that MHC class I haplotypes determined whether

- 39 -

Rhesus monkeys could mount a CTL response to HCV-Core.

Since different MHC class I alleles can bind and present different sets of peptides, these data did not rule out that the Core-ISCOM formulation was not sub-optimal, i.e.

- 5 It remained possible that these animals could mount a Core-specific CTL response directed against peptides other than 86-100 and 121-135. To address this possibility, AY921 and BB231 were challenged with  $2 \times 10^8$  pfu of rVVC/E1 eleven weeks after their 4<sup>th</sup> immunization with Core-ISCOM. As opposed to BB232 and DV036 which had Core-specific CTLs two weeks post rVVC/E1 infection (Table III), neither AY921  
10 nor BB231 had any detectable CTLs post rVVC/E1 inoculation. This strongly suggested that the absence of detectable Core-specific CTLs after Core-ISCOM immunization of these animals was not due to a sub-optimal vaccine formulation, but to an intrinsic inability of these animal to mount such a response, presumably a consequence of their MHC class I haplotype.

15

- To investigate whether immunization with Core-ISCOM induced long-lived CTLs, we monitored DV037 for up to 51 weeks (1 year) after its 4<sup>th</sup> immunization. Peptide 121-135-specific CTLs were detected 10, 15, 31, 38, 45 and 51 weeks post last immunization (Figure 3A). In contrast, the 121-135-specific CTL response primed  
20 by rVVC/E1 in BB232 was barely detectable 14 weeks post vaccination and was undetectable 18 weeks post vaccination (Figure 3B). Similarly, the 86-100-specific CTLs primed in DV036 by rVVC/E1 vaccination became undetectable 14 weeks post vaccination.

- 25 In an effort to quantify the number of peptide 121-135-specific CTLs present in DV037 1 year after it had received its last boost, the animal's PBMCs were restimulated *ex vivo* with 121-135 and the percent of specific CD8<sup>+</sup> T cells was assessed by intracellular staining for IFN- $\gamma$  and TNF- $\alpha$ . As illustrated in Figure 3C (left panels), 121-135-specific CTLs represented 0.49% of the peripheral CD8<sup>+</sup> T  
30 cells (or 490 cells per  $10^5$  CD8<sup>+</sup> T cells) secreted IFN- $\gamma$  and/or TNF- $\alpha$  after *ex vivo* peptide stimulation for 12 hours. In contrast, after *in vitro* restimulation with peptide



- 40 -

121-135, 71% of these CD8+ T cells were specific for this peptide as determined by their abilities to secrete IFN- $\gamma$  and/or TNF- $\alpha$  (Figure 3C, right panels).

#### Example 4

#### 5 Characterization of Cellular and Humoral Immune Responses

##### In Rhesus Monkeys Immunized with Core-ISCOM

Although only one out of three Core-ISCOM immunized animals had detectable CTLs, the fact that in the responding animal Core-specific CTLs were detected after  
10 only two immunizations (Table III) and were long-lived (Figure 3A) formed the basis to immunize five more animals (15860-4) with Core-ISCOM (see Table II for dosage and immunization schedule). All animals were naïve at the time of vaccination. In this study, the priming of both Core-specific CTLs and Core-specific CD4+ T cells and antibodies were monitored.

15 None of the animals had any detectable Core-specific CD4+ or CD8+ T cells at the time of immunization (week 0, Table IV). Core-specific CD4+ T cells, as determined by lymphoproliferation assay, were detected in all animals except 15861 after the second immunization, but this animal had a detectable CD4+ response after the third  
20 immunization (Table IV). For animals 15862 and 15863, it is unlikely that the low SI observed after the 3<sup>rd</sup> immunization (Table IV) was due to the absence of a CD4+ T cell response as a strong proliferation was observed in these animals and at this particular time point for the *E. coli* control (see below). None of the animals had antibodies against Core prior to immunization. However, all animals had  
25 seroconverted to Core after two immunizations, and the level of antibodies against Core was boosted by a third immunization (Figure 4). Notably, the mean Core antibody titer among these animals was comparable after two immunization (1,931) and higher after three immunization (4,566) (Figure 4) to that present in the serum of chronically infected patients with an unusually high anti-Core antibody titer (2,358;  
30 not shown) run in the same assay.

- 41 -

To investigate whether Core-ISCOM elicited a Th1 or Th2-type response in these monkeys, freshly isolated PBMCs prior to vaccination as well as two weeks post 2<sup>nd</sup> and 3<sup>rd</sup> immunizations were tested for their capacity to produce cytokines at 48h in response to stimulation with Core peptides spanning the entire length of the Core protein. As shown in Figures 5A and 5B, a significant increase in Th1 cytokines (IFN- $\gamma$  and IL-2) was observed post immunization in all animals, although the magnitude of response observed for 15860, 15861 and 15862 was lower than that observed for animals 15863 and 15864. Similarly, an increase in Th2-type cytokines (IL-5 and IL-10) was observed in all animals following vaccination, with the highest amount of IL-5 and IL-10 also detected in 15863 and 15864 (Figures 5C and 5D). Although the amount of secreted Th2-type cytokines were lower than that detected for Th1-type cytokines, these data indicated that Core-ISCOM induced a Th0-like type response in Rhesus monkeys.

After two and three immunizations, three animals (15862, 15863 and 15864) had detectable CTL responses directed against pool B (aa: 60-140), while the two others (15860 and 15861) did not (Table IV). This CTL response was directed against peptide 86-100 for animal 15864 and against peptide 121-135 for animals 15862 and 15863. Since 86-100 and 121-135-specific CTLs were also primed in DV036 (86-100), DV037 (121-135) and BB 232 (121-135) (Table III), it was of importance to determined whether all 86-100 and 121-135-specific CTLs were respectively restricted by a single MHC class I allele. The 86-100-specific CTL line derived from animal 15864 efficiently lysed peptide 86-100-sensitized B-LCLs derived from 15864 but not peptide-sensitized B-LCLs derived from animal DV036, indicating that a different (unidentified) MHC class I allele presented this peptide to CTL (Figure 6A

- 42 -

specific CTLs in these animals. Taken together, this suggested, once again, that the MHC class I haplotype of the animals dictated whether they could mount Core-specific CTLs.

- 5 In an effort to quantitate the number of Core-specific CD8+ and CD4+ T cells primed in the animals described above, freshly isolated PBMCs were stained for intracellular INF- $\gamma$  and TNF- $\alpha$  after *ex vivo* restimulation. The CD8+ T cell responses to naturally processed peptides were quantified after *ex vivo* restimulation with autologous B-LCLs infected with rVVC/E1 or VVwt, as a control. The CD4+ T cell responses to
- 10 naturally processed peptides were quantified after *ex vivo* restimulation with the recombinant Core protein or an *E. coli* control. Intracellular staining responses revealed that while none of the animals had detectable Core-specific CD8+ T cells at the time of immunization, between 0.30 and 0.71% of 15862, 15863 and 15864's peripheral CD8+ T cells were specific for naturally processed Core-derived peptide(s)
- 15 after 2 immunizations (Figure 7A). The number of specific CTLs was, however, not increased after the third immunization, as judged by intracellular staining responses. Notably, no CD8+ T cells secreting IFN- $\gamma$  and/or TNF- $\alpha$  in response to Core were detected in the two animals (15860 and 15861) for whom no Core-specific CTL activity was observed by  $^{51}\text{Cr}$  release assay (Figure 7A and Table IV). Quantification
- 20 of Core-specific CD4+ T cells confirmed the data obtained by lymphoproliferation assay (Table IV) in that between 0.32 and 2.21% of CD4+ T cells from all five animals were specific for naturally-processed Core peptides (Figure 7B). Furthermore, the fact that 0.53 and 0.28% of CD4+ T cells from animals 15862 and 15863 were positive for cytokines after the 3<sup>rd</sup> immunization (Figure 7B), strongly
- 25 suggested that the negative SI observed for this time point (Table IV) was indeed a 'false negative', most likely due to the high proliferation observed in response to the *E. coli* control. This suggested that intracellular staining for IFN- $\gamma$  and TNF- $\alpha$  is a more sensitive assay than lymphoproliferation to assess antigen-specific CD4+ T cell responses. Indeed, for animal 15861, no CD4+ T cells were detected by
- 30 lymphoproliferation 2 weeks post 2<sup>nd</sup> immunization (Table IV). In contrast, Core-specific CD4+ T cells were detected in this animal (2 weeks post 2<sup>nd</sup>) by intracellular staining for IFN- $\gamma$  and TNF- $\alpha$  (Figure 7B).

### Example 5

#### The Use of Core-ISCOMs as Adjuvants for HCV Polypeptides

- 5 Because vaccination with recombinant HCV envelope proteins and adjuvant can, at least in some instances, influence the outcome of infection and disease (Choo *et al.* (1994) *Proc. Natl. Acad. Sci. USA* 91:1294), we investigated whether the Core-ISCOM above could serve as an adjuvant for other HCV polypeptides, such as the heterodimeric envelope protein E1E2. To that end, mice (10 animals per group) were
- 10 immunized with 2µg of soluble E1E2 protein alone, or 2µg of soluble E1E2 in the presence of the adjuvant MF59, or in the presence of 2µg of Core-ISCOM. As shown in Figure 8, mice immunized with E1E2 alone had no significant anti-E2 antibody titer. In contrast, mice immunized with E1E2 + Core-ISCOM had a significant anti-E2 antibody titer after three immunizations, and these titers were comparable to those
- 15 observed in mice immunized with E1E2 + MF59. Furthermore, the 'quality' of antibody elicited in mice immunized by E1E2 + MF59 and E1E2 + Core-ISCOM appeared to be comparable with the antibody titers that could inhibit the binding of HCV-1a E2 to the HCV putative receptor CD81 in both groups of mice (Figure 9).
- 20 The above examples demonstrate that vaccination with HCV polypeptide ISCOMs are able to prime strong HCV polypeptide-specific CD8+ and CD4+ T cells as well as anti-HCV polypeptide antibodies. Furthermore these ISCOM formulations are able to serve as adjuvants to elicit antibodies against other HCV proteins. Thus, HCV ISCOMs may prevent the establishment of chronicity, and/or increase the response
- 25 rate to anti-viral therapy.

### Example 6

#### Sucrose Gradient Analysis of NS35Core121-ISCOM

- 30 The NS35Core121 protein was found in a broad peak across the gradient (Figure 10A). The NS35Core121 ISCOM the protein was essentially found in fractions 15 to 20 which corresponded to an ISCOMATRIX™ peak indicating association has

- 44 -

occurred (Figure 10B). Interestingly an ISCOMATRIX™ was also found in fractions 5 to 10 which indicates there was a proportion of the ISCOMATRIX™ with no protein associated.

- 5 Accordingly, novel HCV/ISCOM compositions and methods of using the same have been disclosed. From the foregoing, it will be appreciated that, although
- 
- specific embodiments of the invention have been described herein for purposes of illustration, various modifications may be made without deviating from the spirit and scope of the appended claims.

- 45 -

**Table I - Stability of Core-ISCOM™ Formulations**

Time months	Particle size	Sucrose gradient
0	1.5	>90% associated
1	1.7	>90% associated
11	1.8	>90% associated

**Table II – Summary of Immunization and MHC-I type.**

Animal Number	MHC class I Tping (Mamu A* & B*) <sup>a</sup>	Immunogen	Dose (Route)	Immunization schedule (weeks)
<b>First Study</b>				
BB228 BB232 DV036	A*08; B*17 B*03 B*03	rVVC/E1	1x10 <sup>8</sup> pfu (scarification) 1x10 <sup>8</sup> pfu (ID)	0
AY921 <sup>b</sup> BB231 <sup>b</sup> DV037	A*08; B*03 A*01; A*02 B*03; B*04	HCV Core ISCOM™	25 µg (IM)	0, 4, 8, 27
AY922 BB227 BB230	B*01 A*02 A*08; B*03	HCV Core Adjuvant: LTK63	200 µg (IM) 200 µg (IM)	0, 4, 8, 27
<b>Second Study</b>				
15860 <sup>b</sup> 15861 <sup>b</sup> 15862 15863 15864	B*01; B*03 B*03 B*03 - B*01	HCV Core ISCOM™	50 µg (IM)	0, 4, 8

<sup>a</sup> Mamu alleles tested for: A\*01, \*02, \*08 and 11; B01, \*03, \*04 and \*17.

<sup>b</sup> These animals also received 2 x 10<sup>8</sup> pfu or rVVC/E1 nine to eleven weeks post last immunization.

ID: Intradermal; IM: intramuscular.

Table III – Priming of Core-specific CTLs in Rhesus macaques.

Week	E:T Ratio	Percent Specific Lysis <sup>a</sup>											
		DV037 (Core-ISCMS)			BB230 (Core-LTK63) Core Pool 2 (aa: 41-90)			BB232 (rVVC/E1) Core pool 4 (aa: 121-170)			DV036 (rVVC/E1) Core pool 3 (aa: 81-130)		
		Hm <sup>b</sup>	Hit <sup>b</sup>	Hm <sup>b</sup>	Hm <sup>b</sup>	Hit <sup>b</sup>	Hm <sup>b</sup>	Hm <sup>b</sup>	Hit <sup>b</sup>	Hm <sup>b</sup>	Hm <sup>b</sup>	Hit <sup>b</sup>	Hm <sup>b</sup>
0 (pre)	40:1 13:1 4:1	<1 10 2	<1 <1 <1	11 <1 <1	<1 <1 <1	NT NT NT	9 7 7	<1 <1 <1	<1 <1 <1	8 <1 <1	12 4 <1		
2 (2w post 1 <sup>st</sup> )	40:1 13:1 4:1	NT NT NT	NT NT NT	NT NT NT	NT NT NT	NT NT NT	44 <sup>c</sup> 25 13	<1 <1 <1	<1 <1 <1	24 <sup>d</sup> 15 6	5 <1 <1		
6 (2w post 2 <sup>nd</sup> )	40:1 13:1 4:1	20 10 5	<1 <1 <1	17 <1 <1	6 13 10	NT NT NT	N/A N/A N/A	N/A N/A N/A	N/A N/A N/A	N/A N/A N/A	N/A N/A N/A		
10 (2w post 3 <sup>rd</sup> )	40:1 13:1 4:1	20 12 <1	<1 <1 <1	49 23 14	9 <1 <1	NT NT NT	N/A N/A N/A	N/A N/A N/A	N/A N/A N/A	N/A N/A N/A	N/A N/A N/A		
14 (6w post 3 <sup>rd</sup> )	40:1 13:1 4:1	30 <sup>c</sup> 16 12	5 <1 <1	40 14 9	30 15 10	NT NT NT	N/A N/A N/A	N/A N/A N/A	N/A N/A N/A	N/A N/A N/A	N/A N/A N/A		
29 (2w post 4 <sup>th</sup> )	40:1 13:1 4:1	50 35 28	2 3 2	17 <1 <1	<1 <1 <1	NT NT NT	N/A N/A N/A	N/A N/A N/A	N/A N/A N/A	N/A N/A N/A	N/A N/A N/A		

NT: Not Tested; N/A: Not applicable.

<sup>a</sup> Percent Specific Lysis is only shown for animals with detectable CTL activity and for the peptide pools against which such CTL activity was detected.<sup>b</sup> CT- activity of CD8+ T cells restimulated *in vitro* with a peptide pool (1 through 5) was tested against autologous B-LCLs sensitised with the same peptide pool (Hm) or an irrelevant peptide pool (Hit).<sup>c</sup> The epitopic peptide recognised is aa: 121-135 (KVVDLTTCGFADLMG). <sup>d</sup> The epitopic peptide recognised is aa: 88-100 (YGNEGCGWAGWLLSP).

Table IV – Priming of Core-specific CD8+ and CT4+ in Rhesus macaques immunized with core-ISCOMs.

Week	E:T Ratio	Animal #15860				Animal #15861				Animal #15862				Animal #15863				Animal #15864			
		CD8 resp. (%)		CD4+ resp. (%)		CD8 resp. (%)		CD4+ resp. (%)		CD8 resp. (%)		CD4+ resp. (%)		CD8 resp. (%)		CD4+ resp. (%)		CD8 resp. (%)		CD4+ resp. (%)	
		Lysis) <sup>a</sup>	Core pool B (aa: 61-140)	Lysis) <sup>a</sup>	Core pool B (aa: 61-140)	Lysis) <sup>a</sup>	Core pool B (aa: 61-140)	Lysis) <sup>a</sup>	Core pool B (aa: 61-140)	Lysis) <sup>a</sup>	Core pool B (aa: 61-140)	Lysis) <sup>a</sup>	Core pool B (aa: 61-140)	Lysis) <sup>a</sup>	Core pool B (aa: 61-140)	Lysis) <sup>a</sup>	Core pool B (aa: 61-140)	Lysis) <sup>a</sup>	Core pool B (aa: 61-140)	Lysis) <sup>a</sup>	Core pool B (aa: 61-140)
		Hm	Ht	Hm	Ht	Hm	Ht	Hm	Ht	Hm	Ht	Hm	Ht	Hm	Ht	Hm	Ht	Hm	Ht	Hm	Ht
0	40:1	8	13	4	1	14	14	11	5	11	5	11	5	11	5	11	5	11	5	11	5
	13:1	2	11	<1	<1	6	8	<1	4	<1	4	0.9	1.2	<1	13	<1	13	<1	13	<1	0.8
	4:1	2	5	<1	<1	3	<1	1	3	1	3	1	3	1	<1	<1	<1	<1	<1	<1	<1
6	40:1	8	3	7	<1	31 <sup>c</sup>	8	34 <sup>c</sup>	9	34 <sup>c</sup>	9	34 <sup>c</sup>	9	34 <sup>c</sup>	9	34 <sup>c</sup>	9	34 <sup>c</sup>	9	34 <sup>c</sup>	9
	13:1	<1	2	<1	<1	16	6	16	6	16	6	5.0	3.5	16	6	20	4	20	4	20	4
	4:1	<1	<1	<1	<1	4	<1	8	1	8	1	1	3	8	1	12	3	12	3	12	3
10	40:1	<1	3	13	13	31	5	29	1	29	1	2.1 <sup>e</sup>	1.1 <sup>e</sup>	29	1	79	10	79	10	79	10
	13:1	<1	12	5	7	14	4	17	<1	17	<1	4	<1	17	<1	60	11	60	11	60	11
	4:1	<1	<1	<1	2	6	<1	8	<1	8	<1	<1	<1	8	<1	33	11	33	11	33	11

<sup>a</sup> CTL activity of CD8+ T cells restimulated *in vitro* with a peptide pool (A through C) was tested against autologous B-LCLs sensitized with the same peptide pool (Hm) or an irrelevant peptide pool (H1). Percent Specific Lysis is only shown for pool B since no CTL activity was detected in any animals for pool A (aa: 1-70) or pool C (aa: 131-191).

<sup>b</sup> CD4+ response was determined by lymphoproliferation assay as described in Materials and Methods. A stimulation index (SI) of 3.0 or greater was scored positive.

<sup>c</sup> epitope recognised was aa: 121-130 (KVIDTLTCGFADLMG).

<sup>d</sup> epitope recognised was aa: 86-100 (YGNEGCGWAGWLLSP).

<sup>e</sup> High proliferation to *E. coli* control.



**CLAIMS:**

1. An immunogenic complex comprising a charged organic carrier and a charged antigen, which organic carrier and antigen are electrostatically associated, and wherein the charged antigen is a polyprotein of Hepatitis C Virus (HCV) or a fragment thereof, or a fusion protein comprising said polyprotein or a fragment thereof.
2. The immunogenic complex according to claim 1, wherein said polyprotein is the core protein of HCV.
3. The immunogenic complex according to claim 1, wherein said carrier is negatively charged.
4. The immunogenic complex according to claim 1 wherein said carrier is an adjuvant.
5. The immunogenic complex according to claim 4 wherein said carrier is a negatively charged adjuvant.
6. The immunogenic complex according to claim 5 wherein said negatively charged adjuvant is a naturally negatively charged adjuvant.
7. The immunogenic complex according to claim 5 wherein said negatively charged adjuvant is a naturally negatively charged adjuvant which has been modified to increase the degree of its negative charge.
8. The immunogenic complex according to claim 4 wherein said adjuvant comprises a saponin.
9. The immunogenic complex according to claim 4 wherein said adjuvant is a saponin complex.

10. The immunogenic complex according to claim 9 wherein said saponin complex is ISCOMATRIX™.
11. The immunogenic complex according to claim 4 wherein said adjuvant comprises a phospholipid.
12. The immunogenic complex according to claim 11 wherein said phospholipid is a phosphoglyceride.
13. The immunogenic complex according to claim 12 wherein the phosphoglyceride is selected from the group consisting of phosphatidyl inositol, phosphatidyl glycerol, phosphatidic acid and cardiolipin.
14. The immunogenic complex according to claim 11 wherein said phospholipid is lipid A.
15. The immunogenic complex according to claim 14 wherein the lipid A is selected from the group consisting of diphosphoryl lipid A and monophosphoryl lipid A.
16. The immunogenic complex according to claim 1 wherein said complex induces a cytotoxic T-lymphocyte response.
17. A vaccine composition comprising as the active component an immunogenic complex comprising a charged organic carrier and a charged antigen, which carrier and antigen are electrostatically associated, and wherein the charged antigen is a polyprotein of Hepatitis C Virus (HCV) or a fragment thereof, or a fusion protein comprising said polyprotein or a fragment thereof, together with one or more pharmaceutically acceptable carriers and/or diluents.

- 50 -

18. The vaccine composition according to claim 17 wherein said polyprotein is the core protein of HCV.
  19. The vaccine composition according to claim 17 wherein said carrier is negatively charged.
- 
20. The vaccine composition according to claim 17 wherein said carrier is an adjuvant.
  21. The vaccine composition according to claim 20 wherein said carrier is a negatively charged adjuvant.
  22. The vaccine composition according to claim 21 wherein said negatively charged adjuvant is a naturally negatively charged adjuvant.
  23. The vaccine composition according to claim 21 wherein said negatively charged adjuvant is a naturally negatively charged adjuvant which has been modified to increase the degree of its negative charge.
  24. The vaccine composition according to claim 20 wherein said adjuvant comprises a saponin.
  25. The immunogenic complex according to claim 20 wherein said adjuvant is a saponin complex.
  26. The vaccine composition according to claim 25 wherein said saponin complex is ISCOMATRIX™.
  27. The vaccine composition according to claim 20 wherein said adjuvant comprises a phospholipid.

- 51 -

28. The vaccine composition according to claim 27 wherein said phospholipid is a phosphoglyceride.
  29. The vaccine composition according to claim 28 wherein the phosphoglyceride is selected from the group consisting of phosphatidyl inositol, phosphatidyl glycerol, phosphatidic acid and cardiolipin.
- 
30. The vaccine composition according to claim 27 wherein said phospholipid is lipid A.
  31. The vaccine composition according to claim 30 wherein the lipid A is selected from the group consisting of diphosphoryl lipid A and monophosphoryl lipid A.
  32. The vaccine composition according to claim 17 wherein said composition induces a cytotoxic T-lymphocyte response.
  33. The vaccine composition according to claim 17 further comprising one or more additional HCV protein.
  34. The vaccine composition according to claim 33 wherein said additional HCV protein is selected from the group consisting of a nonstructural protein, the E1 envelope protein, the E2 envelope protein, an immunogenic fragment of one of these proteins, and combinations of these proteins and fragments.
  35. A method of eliciting, inducing or otherwise facilitating, in a mammal, an immune response to an antigen, said method comprising administering to said mammal an effective amount of an immunogenic complex according to claim 1.
  36. A method of eliciting, inducing or otherwise facilitating, in a mammal, an immune response to an antigen, said method comprising administering to said mammal an effective amount of a vaccine composition according to claim 17.




37. The method according to claim 35 or 36 wherein said immune response comprises a cytotoxic T-lymphocyte response.
38. A method of treating a disease condition in a mammal said method comprising ~~administering to said mammal an effective amount of an immunogenic~~ complex according to claim 1 wherein administering said complex elicits, induces or otherwise facilitates an immune response which inhibits, halts, delays or prevents the onset or progression of said disease condition.
39. A method of treating a disease condition in a mammal said method comprising administering to said mammal an effective amount of a vaccine composition according to claim 17 wherein administering said composition elicits, induces or otherwise facilitates an immune response which inhibits, halts, delays or prevents the onset or progression of the disease condition.
40. The method according to claim 38 or 39 wherein said immune response comprises a cytotoxic T-lymphocyte response.
41. The method according to claim 38 or 39 wherein said treatment is therapeutic treatment of said disease condition.
42. The method according to claim 38 or 39 wherein said treatment is prophylactic treatment of said disease condition.
43. The method according to claim 38 or 39 wherein said disease condition results from an HCV infection.
44. Use of an immunogenic complex according to claim 1 in the manufacture of a medicament for inhibiting, halting, delaying or preventing the onset or progression of a disease condition.

- 53 -

- 45. Use of a vaccine composition according to claim 17 in the manufacture of a medicament for inhibiting, halting, delaying or preventing the onset or progression of a disease condition.
  - 46. Use according to claim 44 or 45 wherein said disease condition results from an HCV infection.
- 
- 47. An agent for use in inhibiting, halting, delaying or preventing the onset or progression of a disease condition wherein said agent comprises an immunogenic complex according to claim 1.
  - 48. An agent for use in inhibiting, halting, delaying or preventing the onset or progression of a disease condition wherein said agent comprises a vaccine composition according to claim 17.
  - 49. An agent according to claim 47 or 48 wherein said disease condition results from an HCV infection.


1/10

Figure 1



2/10

Figure 2





3/10

Figure 3

Fig.3A  
DV037  
Core-ISCOM




Fig. 3B  
BB232  
rVVC/E1






Fig. 3C  
DV037  
51 weeks  
post 4th




4/10

Figure 4




5/10

Figure 5



6/10

Figure 6



7/10

Figure 7

Fig. 7A






Fig. 7B



8/10

Figure 8




9/10

Figure 9



10/10

Figure 10





## INTERNATIONAL SEARCH REPORT

International application No.  
**PCT/AU00/01410****A. CLASSIFICATION OF SUBJECT MATTER**Int. Cl. <sup>7</sup>: A61K 39/385, 39/29; A61P 31/12

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)

IPC 7

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

**SEE DATABASES BELOW**

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)


WPAT, CHEMICAL ABSTRACTS, MEDLINE: Keywords used - saponin, quil, iscom, iscomatrix, hepatitis c virus, hcv, organic carrier

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X	AU A 26515/00 (CSL Limited) 24 August 2000. See entire document.	1-49
P, X	AU A 55099/99 (SMITHKLINE BEECHAM BIOLOGICALS S.A.) 17 February 2000. See entire document.	1-49
X	AU A 96238/98 (SMITHKLINE BEECHAM BIOLOGICALS S.A.) 18 March 1999. See entire document.	1-49

☒ Further documents are listed in the continuation of Box C ☒ See patent family annex

<p>* Special categories of cited documents:</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier application or patent but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p>		<p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art</p> <p>"&amp;" document member of the same patent family</p>
---	--	---

Date of the actual completion of the international search <b>16 January 2001</b>	Date of mailing of the international search report <b>22 January 2001</b>
Name and mailing address of the ISA/AU AUSTRALIAN PATENT OFFICE PO BOX 200, WODEN ACT 2606, AUSTRALIA E-mail address: pct@ipaustalia.gov.au Facsimile No.: (02) 6285 3929	Authorized officer  <b>JULIE CAIRNDUFF</b> Telephone No.: (02) 6283 2545

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/AU00/01410

**C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	AU A 11456/99 (SMITHKLINE BEECHAM BIOLOGICALS S.A.) 11 March 1999. See entire document.	1-49
X	AU A 47812/97 (SMITHKLINE BEECHAM BIOLOGICALS S.A.) 16 April 1998. See entire document.	1-49
X	<del>AU A 36444/95 (ISCOTEC AB) 25 April 1996.</del> <del>See entire document.</del>	<del>1-49</del>
Y	BIOCHEMICAL AND BIOPHYSICAL RESEARCH COMMUNICATIONS volume 240, pages 793-797 (1997) by Nakanishi T. et al. "Positively charged liposome functions as an efficient immunoadjuvant in inducing immune responses to soluble proteins". See entire document.	1-49
Y	IMMUNOLOGY AND CELL BIOLOGY volume 76, pages 295-299 (1998) by Morein B. and Bengtsson K.L. "Functional aspects of iscoms". See entire document.	1-49
Y	JOURNAL OF GENERAL VIROLOGY volume 74, pages 1103-1113 (1993) by Selby M.J. et al. "Expression, identification and subcellular localization of the proteins encoded by the hepatitis C viral genome". See entire document.	1-49

**INTERNATIONAL SEARCH REPORT**  
Information on patent family members

International application No.  
**PCT/AU00/01410**

This Annex lists the known "A" publication level patent family members relating to the patent documents cited in the above-mentioned international search report. The Australian Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

Patent Document Cited in Search Report		Patent Family Member			
AU	26515/00	WO	0048630		
AU	55099/00	GB	9817052		
		WO	0007621		
AU	96238/98	EP	1009430		
		GB	9718901		
		WO	9912565		
AU	11456/99	EP	1009382		
		GB	9720982		
		WO	9911241		
AU	47812/97	BR	9711853	NO	991524
		CN	1238696	PL	332633
		CZ	9901167	WO	9815287
		EP	939650	ZA	9708868
		GB	9620795		
AU	36444/95	WO	9611711	NZ	293882
		EP	785802	CA	2201611
		FI	971498		
		NO	971622		
END OF ANNEX					

HPS Trailer Page  
for  
**WEST**

---

UserID: jstucker

Printer: cm1\_8e12\_gbelptr

**Summary**

<u>Document</u>	<u>Pages</u>	<u>Printed</u>	<u>Missed</u>	<u>Copies</u>
WO000137869	67	67	0	1
Total (1)	67	67	0	-

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**